

STA 302 / 1001 H - Summer 2004

Test 1 – June 2, 2004

LAST NAME: _____ FIRST NAME: _____

STUDENT NUMBER: _____

ENROLLED IN: (circle one) STA 302 STA 1001

INSTRUCTIONS:

- Time: 60 minutes
- Aids allowed: calculator.
- A table of values from the t distribution is on the last page (page 7).
- Total points: 40

Some formulae:

$$b_1 = \frac{\sum(X_i - \bar{X})(Y_i - \bar{Y})}{\sum(X_i - \bar{X})^2}$$

$$b_0 = \bar{Y} - b_1\bar{X}$$

$$\text{Var}(b_1) = \frac{\sigma^2}{\sum(X_i - \bar{X})^2}$$

$$\text{Var}(b_0) = \sigma^2 \left(\frac{1}{n} + \frac{\bar{X}^2}{\sum(X_i - \bar{X})^2} \right)$$

$$\text{Cov}(b_0, b_1) = -\frac{\sigma^2\bar{X}}{\sum(X_i - \bar{X})^2}$$

$$\text{SSTO} = \sum(Y_i - \bar{Y})^2$$

$$\text{SSE} = \sum(Y_i - \hat{Y}_i)^2$$

$$\text{SSR} = b_1^2 \sum(X_i - \bar{X})^2 = \sum(\hat{Y}_i - \bar{Y})^2$$

$$\sigma^2\{\hat{Y}_h\} = \text{Var}(\hat{Y}_h) = \sigma^2 \left(\frac{1}{n} + \frac{(X_h - \bar{X})^2}{\sum(X_i - \bar{X})^2} \right) \quad \sigma^2\{\text{pred}\} = \text{Var}(Y_h - \hat{Y}_h) = \sigma^2 \left(1 + \frac{1}{n} + \frac{(X_h - \bar{X})^2}{\sum(X_i - \bar{X})^2} \right)$$

Working-Hotelling coefficient: $W = \sqrt{2F_{2,n-2;1-\alpha}}$

1	2	3 abcd	3 efg

2. (6 points (2 each)) For each of the following statements, say whether it is true or false. Give a brief justification of your answer.

(a) A value of R^2 close to 1 indicates that the linear regression model is a good fit to the data.

(b) The estimate of the error variance, s^2 , is a random variable.

(c) $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2 = 0$

3. Two species of predatory birds, collard flycatchers and tits, compete for nest holes during breeding season. Frequently, dead flycatchers are found in nest boxes occupied by tits. A field study examined whether the risk of mortality to flycatchers is related to the degree of competition between the two bird species for next sites. At each of 14 locations, the following data were collected: the number of flycatchers killed (the response variable labelled `fc_killed`) and the nest box occupancy measured as a percentage (the predictor variable labelled `tit_occ`).

The data and some SAS output are given below. Some numbers from the SAS output have been purposely deleted.

Location	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<code>fc_killed</code>	0	0	0	0	0	1	1	1	1	2	2	3	4	5
<code>tit_occ</code>	24	33	34	43	50	35	35	38	40	31	43	55	57	64

The REG Procedure
Descriptive Statistics

Variable	Sum	Mean	Uncorrected SS	Variance	Standard Deviation
Intercept	14.00000	1.00000	14.00000	0	0
<code>tit_occ</code>	582.00000	41.57143	25844	126.87912	11.26406
<code>fc_killed</code>	20.00000	1.42857	62.00000	2.57143	1.60357

The REG Procedure
Model: MODEL1
Dependent Variable: `fc_killed`

Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	(A)	19.11669	19.11669		
Error	(B)	14.31188	(C)		
Corrected Total	13	33.42857			

Root MSE	1.09209	R-Square	(D)
Dependent Mean	1.42857	Adj R-Sq	0.5362
Coeff Var	76.44618		

Parameter Estimates

Variable	DF	Parameter Estimate	Standard Error	t Value	Pr > t
Intercept	1	-3.04686	1.15533	-2.64	0.0217
<code>tit_occ</code>	1	0.10766	0.02689	(E)	0.0018

Questions pertaining to this output are on the next two pages.

- (a) (5 points) What are the values of the 5 missing numbers (A through E) in the SAS output?
- (b) (5 points) For the analysis of variance F test, state the null and alternative hypotheses, the value of the test statistic, the distribution of the test statistic under the null hypothesis, the p -value as accurately as possible, and an appropriate conclusion.
- (c) (2 points) Estimate the mean change in the number of flycatchers killed when the nest box tit occupancy increases by 10%.
- (d) (3 points) Give a 95% confidence interval for the slope of the line.

- (e) (5 points) Suppose an additional location was later found to have a nest box tit occupancy of 30%. Give a 90% prediction interval for this new value.
- (f) (3 marks) Would a 90% confidence interval for the mean number of flycatchers killed when the tit occupancy is 30% be wider or narrower than your interval in part (e). Explain why the width of the intervals differ. An answer that only points out the differences in the formulae will receive no marks.
- (g) (2 marks) Basing your answer only on the information you have from the data and SAS output that was given, do you have any concerns about the validity of the prediction interval you found in the part (e)? Explain.