

Accurate directional inference for vector parameters

Nancy Reid

February 26, 2016

with Don Fraser, Nicola Sartori, Anthony Davison



Parametric models and likelihood

- model $f(y; \theta)$,
- data $y = (y_1, \dots, y_n)$
- log-likelihood function
- parameter of interest

$$\theta \in \mathbb{R}^p$$

independent observations

$$\ell(\theta; y) = \log f(y; \theta)$$

$$\theta = (\psi, \lambda), \quad \psi \in \mathbb{R}^d$$


- likelihood inference

$$w(\psi) = 2\{\ell(\hat{\psi}, \hat{\lambda}) - \ell(\psi, \hat{\lambda}_\psi)\} \sim \chi_d^2$$

$$w_B(\psi) = \frac{w(\psi)}{1 + B(\psi)} \sim \chi_d^2 \quad O_p(n^{-2})$$

$$B(\psi) = E\{w(\psi)\}/d$$

$$w^*(\psi) = w(\psi) \left\{ 1 - \frac{\log \gamma(\psi)}{w(\psi)} \right\}$$



Skovgaard, 2001

Example: 2×3 contingency table


- activity amongst psychiatric patients

Everitt, 1992

	Affective disorders	Schizophrenics	Neurotics
Retarded	12	13	5
Not retarded	18	17	25


- model: log-linear $y \sim \text{Poisson}$, $\log\{E(y)\} = X\theta$, $\theta \in \mathbb{R}^6$
- log-likelihood $\ell(\theta; y) = \theta'X'y - 1'e^{X\theta} = \theta's - c(\theta)$
- $\theta = (\psi, \lambda)$ $\psi \in \mathbb{R}^2, \lambda \in \mathbb{R}^4$
- (ψ_1, ψ_2) interaction parameters
- $H_0 : \psi = \psi_0 = (0, 0)$ independence
- log-likelihood $\ell(\psi, \lambda; y) = \psi's_1 + \lambda's_2 - c(\psi, \lambda)$

Testing $\psi = 0$



$$w(\psi_0) \sim \chi_2^2 \quad p\text{-value } 0.047$$

$$w^*(\psi_0) \quad p\text{-value } 0.048$$



$$\text{directional} \quad p\text{-value } 0.050$$

$$\text{exact} \quad p\text{-value } 0.051$$

Linear exponential families

- model $f(y; \theta) = \exp\{\varphi(\theta)^T u(y) - c\{\varphi(\theta)\} - d(y)\}$ y_1, \dots, y_n i.i.d
- sufficient statistic

$$f(s; \theta) = \int_{\{y: s(y)=s\}} f(y; \theta) dy = \exp\{\varphi(\theta)^T s - nc\{\varphi(\theta)\} - \tilde{d}(s)\}$$
- reduce dimension from n to p by marginalization
- linear parameter of interest $\varphi(\theta) = \theta = (\psi, \lambda)$
- model $f(s_1, s_2; \psi, \lambda) = \exp\{\psi^T s_1 + \lambda^T s_2 - nc(\psi, \lambda) - \tilde{d}(s)\}$
- conditional density $f(s_1 | s_2; \psi) = \exp\{\psi^T s_1 - n\tilde{c}_2(\psi) - \tilde{d}_2(s_1)\}$
- reduce dimension from p to d by conditioning

... conditional density

- $f(s_1 | s_2; \psi) = \exp\{\psi^T s_1 - n\tilde{c}_2(\psi) - \tilde{d}_2(s_1)\}$

- $f(s_1 | s_2; \psi) = \frac{f(s_1, s_2; \psi, \lambda)}{f(s_2; \psi, \lambda)} \propto f(s; \psi, \lambda)$ with s_2 held fixed

- s_2 held fixed $\iff \hat{\lambda}_\psi$ held fixed $(\hat{\theta}_\psi = (\psi, \hat{\lambda}_\psi))$

- saddlepoint approximation:

$$f(s_1 | s_2; \psi) \doteq c \exp[\ell(\hat{\theta}_\psi^0; s) - \ell\{\hat{\theta}(s); s\}] |j\{\hat{\theta}(s); s\}|^{-1/2}, \quad s \in L_\psi^0$$

- centering $\ell(\theta; s) = \psi^T s_1 + \lambda^T s_2 + \ell(\theta; y^0)$ fix $s^0 = s(y^0) = 0$

- plane $L_\psi^0 = \{s \in \mathbb{R}^p : s_2 = 0\} = \{s \in \mathbb{R}^p : \hat{\lambda}_\psi = \hat{\lambda}_\psi^0\}$

- $\hat{\theta}(s) : \partial \ell(\theta; s) / \partial \theta = 0$ m.l.e. as a function of the variable

... conditional density

$$f(s_1 | s_2; \psi) \doteq c \exp[\ell(\hat{\theta}_\psi^0; s) - \ell\{\hat{\theta}(s); s\}] |j\{\hat{\theta}(s); s\}|^{-1/2}, \quad s \in L_\psi^0$$



	Affective disorders	Schizophrenics	Neurotics
Retarded	12	13	5
Not retarded	18	17	25

$$s_1 \in \mathbb{R}^2, s_2 \in \mathbb{R}^4$$

L_ψ^0 : all 2×3 tables with the same row and column totals



Directional tests

- measure the **directed departure** from H_0 in L_{ψ}^0
- s_{ψ} : expected value of s_1 , under H_0
- s^0 : observed value of s_1 = 0 from centering
- L_{ψ}^* : line through these two points
- $L_{\psi}^* = ts^0 + (1 - t)(s_{\psi} - s^0), \quad t \in \mathbb{R}$



... directional tests

$$L_{\psi}^* = ts^0 + (1 - t)s_{\psi}$$



○ null hypothesis of independence $t = 0$

X observed value of s $t = 1$

p -value compares probability from X to ∞ to that from ○ to ∞
 along the line in the sample space curve in the parameter space

like a 2-sided p -value $\Pr(\text{response} > \text{observed} \mid \text{response} > 0)$

... directional p -value

- $p\text{-value} = \frac{\int_1^\infty t^{d-1} f\{s(t); \psi\} dt}{\int_0^\infty t^{d-1} f\{s(t); \psi\} dt}$ $s(t)$ along the line L_ψ^*

- t^{d-1} from change to polar coordinates $\|s\|$, conditional on $s/\|s\|$

- Simplifications:


- 1: $L_\psi^* \subset L_\psi^0 \subset \mathbb{R}^p$

- 2: ratio of two integrals – drop any terms that don't depend on t

- 3: saddlepoint approximation

$$f(s; \psi) \doteq c \exp[\ell(\hat{\theta}_\psi^0; s) - \ell\{\hat{\theta}(s); s\}] |j\{\hat{\theta}(s); s\}|^{-1/2}, \quad s \in L_\psi^0$$

2×3 table



$$t = 0$$

10	10	10
20	20	20

$$t = 0.5$$


11.0	11.5	7.5
19.0	18.5	22.5





$$t = 1$$

12	13	5
18	17	25

$$t = 2$$

14	16	0
16	14	30




... 2×3 table

... 2×3 table

simulations

Nominal	0.010	0.025	0.050	0.100	0.250	0.500
LRT	0.011	0.028	0.055	0.107	0.260	0.510
Directional	0.010	0.024	0.050	0.100	0.250	0.501
Skovgaard, 2001	0.010	0.025	0.050	0.101	0.251	0.501
Nominal	0.750	0.900	0.950	0.975	0.990	
LRT	0.757	0.905	0.952	0.974	0.992	
Directional	0.752	0.902	0.950	0.973	0.992	
Skovgaard, 2001	0.752	0.900	0.950	0.973	0.991	

Example: comparison of normal variances







Likelihood ratio statistic	0.0042
Directional	0.0389
Skovgaard, 2001	0.0622
Bartlett's test	0.0136

F-test





Simulations

3 groups, 10 observations per group







Simulations

3 groups, 5 observations per group



Simulations

1000 groups, 5 observations per group



dimension ψ 999; dimension nuisance 1001

Example: covariance selection

- model $y_i \sim N_q(\mu, \Lambda^{-1})$

inverse covariance matrix

- linear exponential family

$$\ell(\theta; y) = \frac{n}{2} \log |\Lambda| - \frac{1}{2} \text{tr}(\Lambda y^T y) + \mathbf{1}^T y \xi - \frac{n}{2} \xi^T \Lambda \xi \quad s_1, s_2$$

- $\theta = (\xi, \Lambda) = (\Lambda \mu, \Lambda)$

- H_0 : some off-diagonal elements of Λ are 0 $\psi_1 = \dots = \psi_d = 0$
conditional independence

- need constrained m.l.e: use `fitConGraph` in `ggm`

- $\hat{\Lambda}^{-1}(t) = t\hat{\Lambda}^{-1} + (1-t)\hat{\Lambda}_0^{-1}$ m.l.e. along the line

- $f\{s(t); \psi\} \propto |t\hat{\Lambda}^{-1} + (1-t)\hat{\Lambda}_0^{-1}|^{(n-q-2)/2}$

... covariance selection

simulations

Nominal	0.010	0.025	0.050	0.100	0.250	0.500
LRT	0.055	0.105	0.170	0.270	0.487	0.730
Directional	0.011	0.026	0.050	0.101	0.248	0.498
Skovgaard, 2001	0.007	0.018	0.036	0.074	0.196	0.422
Nominal	0.750	0.900	0.950	0.975	0.990	
LRT	0.895	0.967	0.985	0.994	0.998	
Directional	0.749	0.899	0.949	0.974	0.990	
Skovgaard, 2001	0.678	0.852	0.919	0.955	0.980	

Covariance matrix 11×11 ; dimension of $\psi = 45$

first order Markov dependence

Tangent exponential model $n \downarrow p$ (dimension of y to dimension of θ)

- Every model $f(y; \theta)$ on \mathbb{R}^n can be approximated by an exponential family model:

$$f_{TEM}(s; \theta) ds = \exp\{\varphi(\theta)'s + \ell(\theta)\} h(s) ds$$

- s is a score variable on \mathbb{R}^p $s(y) = -\ell_\varphi(\hat{\theta}^0; y)$
- $\ell(\theta) = \ell(\theta; y^0)$ is the observed log-likelihood function
- $\varphi(\theta) = \varphi(\theta; y^0)$ is the canonical parameter $\in \mathbb{R}^p$ to be described
- matches log-likelihood function at y^0 , and its first derivative on the sample space, at y^0 by construction
- implements conditioning on an approximate ancillary statistic contrast with exp fam

Aside: canonical parameter $\varphi(\theta)$

- if $f(y; \theta)$ is an exponential family, φ is sitting in the model
- if not find a **pivotal quantity** $z_i = z_i(y_i; \theta)$ with a fixed distribution
example: $(y_i - \mu)/\sigma$

- define $V_i = - \left(\frac{\partial z_i}{\partial y_i} \right)^{-1} \frac{\partial z_i}{\partial \theta} \Big|_{y=y^0, \theta=\hat{\theta}^0}$ a vector of length p

$$\varphi(\theta) = \varphi(\theta; y^0) = \sum_{i=1}^n \frac{\partial \ell(\theta; y^0)}{\partial y_i} V_i = \ell_{;V}(\theta; y^0)$$

Testing a value for ψ

eliminating nuisance parameter: $p \downarrow d$

- $\theta = (\psi, \lambda)$; $\varphi = \varphi(\theta)$ is the canonical parameter of the TEM
- with ψ fixed by the hypothesis, we can integrate out the nuisance parameter by **Laplace approximation**
- define the same plane in the score space L_ψ^0 , where $\hat{\varphi}_\psi$ is fixed at its observed value

$$f(s; \psi) = c \exp\{\ell(\hat{\varphi}_\psi^0; s) - \ell(\hat{\varphi}(s); s)\} |j_{\varphi\varphi}(\hat{\varphi}(s); s)|^{-1/2} |j_{(\lambda\lambda)}(\hat{\varphi}_\psi^0; s)|^{1/2},$$

$$s \in L_\psi^0$$

$$p\text{-value} = \frac{\int_1^\infty t^{d-1} f\{s(t); \psi\} dt}{\int_0^\infty t^{d-1} f\{s(t); \psi\} dt}$$

$s(t)$ along the line L_ψ^*

Example: marginal independence

- $y_i \sim N_q(\mu, \Sigma)$, H_0 : some entries of Σ are 0
- estimate Σ under H_0 using `fitCovGraph`
- $\ell(\Sigma) = \frac{n-1}{2} [\log |\varphi(\Sigma)| - \frac{n-1}{2} \text{tr} \{ \varphi(\Sigma) S \}]$, $\varphi(\Sigma) = \Sigma^{-1}$
 $S =$ sample covariance
- $f\{s(t); \psi\} \propto |t\hat{\Sigma} + (1-t)\hat{\Sigma}_0|^{(n-q-2)/2} |j_{(\lambda\lambda)}\{\hat{\Sigma}_0; s(t)\}|^{1/2}$
- $j_{(\lambda\lambda)}$ needs $\partial\ell(\Sigma)/\partial(\sigma_{jk})$ for the non-zero elements
- $j_{\lambda_j\lambda_k}\{\hat{\Sigma}_0; s(t)\} = \frac{n-1}{2} \left(\text{tr}(A_{kj} + t \left[\text{tr}\{(A_{kj} + A_{jk})(\hat{\Sigma}_0^{-1}\hat{\Sigma} - I_q)\} \right]) \right)$
- $A_{kj} = \Sigma_0^{-1}(\partial\Sigma/\partial\lambda_k)\Sigma_0^{-1}(\partial\Sigma/\partial\lambda_j)$

... independence


simulations; $n = 60; 600, d = 1000$

Nominal	0.010	0.025	0.050	0.100	0.250	0.500
LRT	1.00	1.00	1.00	1.00	1.00	1.00
Directional	0.007	0.024	0.049	0.097	0.250	0.497
Skovgaard, 2001	0.000	0.000	0.000	0.001	0.006	0.026
Nominal	0.750	0.900	0.950	0.975	0.990	
LRT	1.00	1.00	1.00	1.00	1.00	
Directional	0.749	0.904	0.951	0.975	0.990	
Skovgaard, 2001	0.099	0.225	0.333	0.440	0.570	
Nominal	0.010	0.025	0.050	0.100	0.250	0.500
LRT	0.065	0.122	0.199	0.311	0.538	0.786
Directional	0.010	0.024	0.049	0.098	0.251	0.496
Skovgaard, 2001	0.003	0.010	0.022	0.050	0.150	0.352
Nominal	0.750	0.900	0.950	0.975	0.990	
LRT	0.927	0.979	0.991	0.997	0.999	
Directional	0.751	0.898	0.950	0.975	0.989	
Skovgaard, 2001	0.622	0.819	0.894	0.943	0.974	


Conclusion

- different way to assess vector parameters
- incorporates information in the direction of departure
- easy to compute: two model fits, plus 1-d numerical integration
- accurate conditionally, by construction, and unconditionally
simulations
- can be used in models of practical interest
- exponential family model not necessary – easily generalized using approximate exponential family model


... conclusion



... conclusion



... conclusion



References

Fraser, D.A.S. and Massam, H. (1985). Conical tests. *Stat. Hefte*

Skovgaard, I.M. (1988). Saddlepoint expansions for directional test probabilities. *JRSS B*

Skovgaard, I.M. (2001). Likelihood asymptotics. *SJS*

Cheah, P.K., Fraser, D.A.S., Reid, N. (1994). Multiparameter testing in exponential models. *Biometrika*

Davison, A.C., Fraser, D.A.S., Reid, N., Sartori, N. (2014). Accurate directional inference for vector parameters in linear exponential families. *JASA*

Fraser, D.A.S., Reid, N., Sartori, N. (2016). Accurate directional inference for vector parameters. submitted