

## Homework 4

due December 6

1. CH 4.7 Assume  $y_1, \dots, y_n$  is an independent sample from the exponential distribution with density  $\theta \exp(-\theta y), y \geq 0, \theta > 0$ .
  - (a) Obtain the uniformly most powerful test of  $H_0 : \theta = \theta_0$  against alternatives  $\theta < \theta_0$ , and derive the power function of the test.
  - (b) To reduce dependence on possible outliers, it is proposed to reject the largest observation. Derive the uniformly most powerful test from the remaining observations, and show that the loss of power corresponds exactly to the replacement of  $n$  by  $n - 1$ .  
The *Renyi representation* of the order statistics  $(Y_{(1)}, \dots, Y_{(n)})$  is

$$Y_{(r)} \stackrel{d}{=} \frac{1}{\theta} \sum_{j=1}^r \frac{E_j}{n+1-j},$$

where  $E_1, \dots, E_n$  are independent exponential random variables with rate 1. (SM, §2.3).

I took this from CH, exercise 4.7. It was revised in the CH solutions book, to read as follows: “To reduce dependence on possible outliers, it is proposed to replace the largest observation by the 2nd largest. Show that the loss of power corresponds exactly to the replacement of  $n$  by  $n - 1$ .”

You do not need to derive the most powerful test based on  $Y_{(1)}, \dots, Y_{(n)}$ , you should simply replace the test statistic  $\sum Y_i$  from (a) by  $t(Y) = \sum^{n-1} Y_{(i)}$ . You should then be able to use the Renyi representation to find the distribution of  $t(Y)$ . To get the “loss of power” result stated in the question, it seems you will need to use  $t'(Y) = t(Y) + Y_{(n-1)}$ , and if that is easier, fine.

2. The  $p$ -value function is not the same as the power function: the latter depends on both the null value and the alternative value(s) of the parameter. From the Neyman-Pearson lemma, we know that for  $H_0 : \theta = \theta_0$  vs.  $H_A : \theta = \theta_A$ , the most powerful test based on a sample  $y = (y_1, \dots, y_n)$  from  $f(y; \theta)$  is based on the likelihood ratio

$$\frac{L(\theta_A; y)}{L(\theta_0; y)}.$$

- (a) For  $\theta$  a scalar parameter, let  $\theta_A = \theta_0 + \delta$ , for fixed  $\delta > 0$ . Use Taylor series expansions to show that the locally most powerful test, where ‘local’ means  $\delta$  near 0, is a function of  $U(\theta_0; y)$ , the score statistic. [slightly amended](#)

- (b) Compare the local power of the test based on  $U$  to the power of the test based on the likelihood root  $r = \pm\sqrt{2\{\ell(\hat{\theta}) - \ell(\theta)\}}$ , for independent, identically distributed observations from the Cauchy distribution with location parameter  $\theta$ :  $f(y_i; \theta) = [\pi\{1 + (y_i - \theta)^2\}]^{-1}$ .

This can only be done numerically, as far as I can tell. For example, simulate a sample of size 10 from the standard Cauchy with `rcauchy(10)`.

### UMP Invariant tests

A composite null hypothesis  $H_0 : \theta \in \Theta_0$  does not specify the distribution of  $Y$  under  $H_0$ , so some means of reduction to a test statistic with a known distribution under  $H_0$  is needed. In some classes of models a group of transformations can be found which leaves both the null and alternative hypotheses unchanged, and a statistic exists which is a maximal invariant for the group. The resulting test, constructed using the Neyman-Pearson lemma on the marginal density of this invariant, is called most powerful invariant. Some notes on the classical theory of testing are posted, see also CH Ch. 5.2,3 and SM Ch. 7.3.3. The usual textbook example of this is the  $t$ -statistic for comparing two normal means with common unknown variance, but many tests in multivariate analysis are also motivated by considerations of invariance.

The next question is based on CH Example 5.17.

3. Assume  $(x_1, y_1), \dots, (x_n, y_n)$  are independent bivariate normal distributions with mean vector  $(\mu_1, \mu_2)$  and covariance matrix the identity. Suppose we are interested in testing  $H_0 : \mu_1^2 + \mu_2^2 = 1$ , against the alternative  $H_1 : \mu_1^2 + \mu_2^2 > 1$ .
- (a) Show  $H_0$  and  $H_1$  are invariant under the rotation group on the parameter space, which has elements

$$g \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} = \begin{bmatrix} \sin a & \cos a \\ \cos a & -\sin a \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \quad 0 < a \leq 2\pi.$$

- (b) Show that  $t = \bar{x}^2 + \bar{y}^2$  is a maximal invariant statistic under the rotation group on the sample space of  $(\bar{x}, \bar{y})$ , the minimal sufficient statistic, that corresponds to the group on the parameter space, and that the distribution of  $t$  depends only on  $\phi = \mu_1^2 + \mu_2^2$ .
- (c) Find the most powerful test of  $H_0$  based on the marginal distribution of  $t$ , and describe the rejection region  $\mathcal{R}$  for a test of size  $\alpha$ .
- (d) (not required, but potentially of interest): In the more usual case where the variance is  $\sigma^2$  and unknown, the maximal invariant is

$$t' = \frac{t}{\Sigma(x_i - \bar{x})^2 + \Sigma(y_i - \bar{y})^2},$$

and  $u = n(2n - 2)t$  follows a non-central  $F$ -distribution with 2 and  $n - 2$  degrees of freedom, and non centrality parameter  $n\phi/\sigma^2$ . In time series

this is the Schuster test for the amplitude of a given component of the periodogram.