

Homework 4

due December 6

1. CH 4.7 Assume y_1, \dots, y_n is an independent sample from the exponential distribution with density $\theta \exp(-\theta y), y \geq 0, \theta > 0$.

- (a) Obtain the uniformly most powerful test of $H_0 : \theta = \theta_0$ against alternatives $\theta < \theta_0$, and derive the power function of the test.
- (b) To reduce dependence on possible outliers, it is proposed to reject the largest observation. Derive the uniformly most powerful test from the remaining observations, and show that the loss of power corresponds exactly to the replacement of n by $n - 1$.

The *Renyi representation* of the order statistics $(Y_{(1)}, \dots, Y_{(n)})$ is

$$Y_{(r)} \stackrel{d}{=} \frac{1}{\theta} \sum_{j=1}^r \frac{E_j}{n+1-j},$$

where E_1, \dots, E_n are independent exponential random variables with rate 1. (SM, §2.3).

2. The p -value function is not the same as the power function: the latter depends on both the null value and the alternative value(s) of the parameter. From the Neyman-Pearson lemma, we know that for $H_0 : \theta = \theta_0$ vs. $H_A : \theta = \theta_A$, the most powerful test based on a sample $y = (y_1, \dots, y_n)$ from $f(y; \theta)$ is based on the likelihood ratio

$$\frac{L(\theta_A; y)}{L(\theta_0; y)}.$$

- (a) For θ a scalar parameter, let $\theta_A = \theta_0 + \delta/\sqrt{n}$, for fixed $\delta > 0$. Use Taylor series expansions to show that the locally most powerful test is a function of $U(\theta_0; y)$, the score statistic.
- (b) Compare the local power of the test based on U to the power of the test based on the likelihood root, for independent, identically distributed observations from the Cauchy distribution with location parameter θ : $f(y_i; \theta) = [\pi\{1 + (y_i - \theta)^2\}]^{-1}$.

UMP Invariant tests

A composite null hypothesis $H_0 : \theta \in \Theta_0$ does not specify the distribution of Y under H_0 , so some means of reduction to a test statistic with a known distribution under H_0 is needed. In some classes of models a group of transformations can be found which leaves both the null and alternative hypotheses unchanged, and a statistic exists which is a maximal invariant for the group. The resulting test, constructed using the Neyman-Pearson lemma on the marginal density of this invariant,

is called most powerful invariant. Some notes on the classical theory of testing are posted, see also CH Ch. 5.2,3 and SM Ch. 7.3.3. The usual textbook example of this is the t -statistic for comparing two normal means with common unknown variance, but many tests in multivariate analysis are also motivated by considerations of invariance.

The next question is based on CH Example 5.17.

3. Assume $(x_1, y_1), \dots, (x_n, y_n)$ are independent bivariate normal distributions with mean vector (μ_1, μ_2) and covariance matrix the identity. Suppose we are interested in testing $H_0 : \mu_1^2 + \mu_2^2 = 1$, against the alternative $H_1 : \mu_1^2 + \mu_2^2 > 1$.

- (a) Show H_0 and H_1 are invariant under the rotation group on the parameter space, which has elements

$$g \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix} = \begin{bmatrix} \sin a & \cos a \\ \cos a & -\sin a \end{bmatrix} \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}, \quad 0 < a \leq 2\pi.$$

- (b) Show that $t = \bar{x}^2 + \bar{y}^2$ is a maximal invariant statistic under the rotation group on the sample space that corresponds to the group on the parameter space, and that the distribution of t depends only on $\phi = \mu_1^2 + \mu_2^2$.
- (c) Find the most powerful test of H_0 based on the marginal distribution of t , and describe the rejection region \mathcal{R} for a test of size α .
- (d) (not required, but potentially of interest): In the more usual case where the variance is σ^2 and unknown, the maximal invariant is

$$t' = \frac{t}{\Sigma(x_i - \bar{x})^2 + \Sigma(y_i - \bar{y})^2},$$

and $u = n(2n - 2)t$ follows a non-central F -distribution with 2 and $n - 2$ degrees of freedom, and non centrality parameter $n\phi/\sigma^2$. In time series this is the Schuster test for the amplitude of a given component of the periodogram.