



**Harris Recurrence of Metropolis-Within-Gibbs  
and Trans-Dimensional Markov Chains**

by

**Gareth O. Roberts**  
Department of Mathematics and Statistics  
Lancaster University

and

**Jeffrey S. Rosenthal**  
Department of Statistics  
University of Toronto

**Technical Report No. 0412 December 1, 2004**

TECHNICAL REPORT SERIES

University of Toronto  
Department of Statistics

# Harris Recurrence of Metropolis-Within-Gibbs and Trans-Dimensional Markov Chains

by

Gareth O. Roberts\* and Jeffrey S. Rosenthal\*\*

(December 2004; revised May 2006)

**Abstract.** A Markov chain with stationary probability distribution, which is  $\phi$ -irreducible and aperiodic, will converge to its stationary distribution from almost all starting points. The property of Harris recurrence allows us to replace “almost all” by “all”, which is potentially important when running Markov chain Monte Carlo algorithms. Full-dimensional Metropolis-Hastings algorithms are known to be Harris recurrent. In this paper, we consider conditions under which Metropolis-Within-Gibbs and transdimensional Markov chains are or are not Harris recurrent. We present a simple but natural two-dimensional counterexample showing how Harris recurrence can fail, and also a variety of positive results which guarantee Harris recurrence. We also present some open problems. We close with a discussion of the practical implications to MCMC algorithms.

## 1. Introduction.

Harris recurrence is a concept introduced fifty years ago by Harris (1956). More recently, connections between Harris recurrence and Markov chain Monte Carlo (MCMC) algorithms were investigated by Tierney (1994) and Chan and Geyer (1994). In this paper, we re-examine Harris recurrence of various MCMC algorithms, in a more general context.

Markov chains with stationary distributions are the basis of MCMC algorithms. For the algorithm to be valid, it is crucial that the chain will converge to stationarity in

---

\* Department of Mathematics and Statistics, Fylde College, Lancaster University, Lancaster, LA1 4YF, England. Internet: [g.o.robert@lancaster.ac.uk](mailto:g.o.robert@lancaster.ac.uk).

\*\* Department of Statistics, University of Toronto, Toronto, Ontario, Canada M5S 3G3. Internet: [jeff@math.toronto.edu](mailto:jeff@math.toronto.edu). Supported in part by NSERC of Canada.

distribution. If the state space is countable, and the Markov chain is aperiodic and also (*classically*) *irreducible* (i.e., has positive probability of reaching any state from any other state), then it is well known that convergence to stationarity is guaranteed from all starting states (see e.g. Hoel et al., 1972; Billingsley, 1995; Rosenthal, 2000).

On the other hand, classical irreducibility is unachievable when the state space  $\mathcal{X}$  is uncountable. A weaker property is  *$\phi$ -irreducibility* (i.e., having positive probability of reaching every subset  $A$  with  $\phi(A) > 0$  from every state  $x \in \mathcal{X}$ , for some non-zero measure  $\phi(\cdot)$ ). It is known that a  $\phi$ -irreducible, aperiodic Markov chain, with stationary probability distribution  $\pi(\cdot)$ , must still converge to  $\pi(\cdot)$  from  $\pi$ -almost every starting point (see e.g. Nummelin, 1984; Meyn and Tweedie, 1993; Rosenthal, 2001). However, if a chain is  $\phi$ -irreducible but not classically irreducible, then it is indeed possible to have a null set of states from which convergence does not occur.

Tierney (1994) and Chan and Geyer (1994) note that this null set of points from which convergence fails could cause practical problems for MCMC algorithms if the user happens to choose an initial state in this null set. Thus, understanding the nature of this null set is important for applications of MCMC, as well as theoretically. Chan and Geyer (1994) refer to this null set as a “measure-theoretic pathology”. However, we shall see herein that the null set can arise quite naturally, on both discrete and continuous state spaces, including for a simple two-dimensional Metropolis-within-Gibbs algorithm with continuous densities.

This paper is structured as follows. Section 2 presents some background about Markov chains and Harris recurrence, and Theorem 6 proves a number of equivalences of Harris recurrence. Section 3 discusses full-dimensional Metropolis-Hastings algorithms, and Section 4 then discusses Metropolis-within-Gibbs algorithms. Example 9 demonstrates that a simple two-dimensional Metropolis-within-Gibbs algorithm, with continuous target and proposal densities, although irreducible and aperiodic, can still fail to be Harris recurrent or to converge to stationarity from all starting points. Sections 4 and 5 prove various positive results which guarantee Harris recurrence for Metropolis-within-Gibbs algorithms under various conditions, and Section 6 does the same for transdimensional Markov chains.

## 2. Markov Chains and Harris Recurrence.

Consider a Markov chain  $\{X_n\}$  with transition probabilities  $P(x, \cdot)$ , on a state space  $\mathcal{X}$  with  $\sigma$ -algebra  $\mathcal{F}$ . Let  $P^n(x, \cdot)$  be the  $n$ -step transition kernel, and for  $A \in \mathcal{F}$ , let  $\tau_A = \inf\{n \geq 1 : X_n \in A\}$  be the first return time to  $A$ , with  $\tau_A = \infty$  if the chain never returns to  $A$ .

Recall that a Markov chain is  $\phi$ -irreducible if there exists a non-zero  $\sigma$ -finite measure  $\psi(\cdot)$  on  $(\mathcal{X}, \mathcal{F})$  such that  $\mathbf{P}[\tau_A < \infty | X_0 = x] > 0$  for all  $x \in \mathcal{X}$  and all  $A \in \mathcal{F}$  with  $\psi(A) > 0$ . The probability distribution  $\pi(\cdot)$  on  $(\mathcal{X}, \mathcal{F})$  is *stationary* for the chain if  $\int_{\mathcal{X}} \pi(dx) P(x, A) = \pi(A)$  for all  $A \in \mathcal{F}$ .

The *period* of a  $\phi$ -irreducible chain with stationary distribution  $\pi(\cdot)$  is the largest  $D \in \mathbf{N}$  (the set of all positive integers) for which there exist disjoint subsets  $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_D \in \mathcal{F}$  with  $\pi(\mathcal{X}_i) > 0$ , such that  $P(x, \mathcal{X}_{i+1}) = 1$  for all  $x \in \mathcal{X}_i$  ( $1 \leq i \leq D-1$ ), and  $P(x, \mathcal{X}_1) = 1$  for all  $x \in \mathcal{X}_D$ . If  $D = 1$ , then the chain is *aperiodic*.

In terms of these definitions, we have the following classical result, as in e.g. Tierney (1994, p. 1758) or Rosenthal (2001). (See also Nummelin, 1984 and Meyn and Tweedie, 1993.)

**Proposition 1.** *Consider a  $\phi$ -irreducible, aperiodic Markov chain with stationary probability distribution  $\pi(\cdot)$ . Let  $G$  be the set of  $x \in \mathcal{X}$  such that  $\lim_{n \rightarrow \infty} \|P^n(x, \cdot) - \pi(\cdot)\| = 0$ . Then  $\pi(G) = 1$ .*

We also note that aperiodicity is not essential in Proposition 1:

**Proposition 2.** *Consider a  $\phi$ -irreducible Markov chain with stationary probability distribution  $\pi(\cdot)$ , and period  $D \geq 1$ . Let  $G$  be the set of  $x \in \mathcal{X}$  such that  $\lim_{n \rightarrow \infty} \|(1/D) \sum_{r=1}^D P^{nD+r}(x, \cdot) - \pi(\cdot)\| = 0$ . Then  $\pi(G) = 1$ .*

**Proof.** If  $D = 1$ , then this reduces to Proposition 1 above. If  $D > 1$ , then let  $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_D$  be as in the definition of period. Then  $P^D$  is  $\phi$ -irreducible and aperiodic when restricted to each  $\mathcal{X}_i$ , with stationary distribution  $\pi_i(\cdot)$ , such that  $\pi(\cdot) = (1/D) \sum_{r=1}^D \pi_r(\cdot)$ . It follows from Proposition 1 that for  $\pi_D$ -a.e.  $x \in \mathcal{X}_D$ , and for any  $1 \leq r \leq D$ , we have  $\lim_{n \rightarrow \infty} \|P^{nD+r}(x, \cdot) - \pi_r(\cdot)\| = 0$ . Hence, for  $\pi_i$ -a.e.  $x \in \mathcal{X}_i$ , we have

$\lim_{n \rightarrow \infty} \|P^{nD+r+D-i}(x, \cdot) - \pi_r(\cdot)\| = 0$ . The result follows by averaging over  $1 \leq r \leq D$  and using the triangle inequality.  $\blacksquare$

The above conclusions still allow for a null set  $G^C$  from which convergence fails. This null set can indeed arise, even for simple examples, on both discrete and continuous state spaces:

**Example 3.** (C. Geyer, personal communication; Roberts and Rosenthal, 2004) Let  $\mathcal{X} = \{1, 2, \dots\}$ . Let  $P(1, \{1\}) = 1$ , and for  $x \geq 2$ ,  $P(x, \{1\}) = 1/x^2$  and  $P(x, \{x+1\}) = 1 - (1/x^2)$ . This chain has stationary distribution  $\pi(\cdot) = \delta_1(\cdot)$ , and it is  $\phi$ -irreducible (with respect to  $\pi$ ) and aperiodic. On the other hand, if  $X_0 = x \geq 2$ , then  $\mathbf{P}[X_n = x+n \text{ for all } n] = \prod_{j=x}^{\infty} (1 - (1/j^2)) > 0$ , so that  $\|P^n(x, \cdot) - \pi(\cdot)\| \not\rightarrow 0$ . Hence, convergence holds only from the set  $G = \{1\}$ , but fails to hold from the set  $G^C = \{2, 3, 4, \dots\}$ . Of course, this example is not irreducible in the classical sense, since no state  $x \geq 2$  is reachable from the state 1. However, it is still *indecomposable* (see e.g. Rosenthal, 1995b).

**Example 4.** (continuous state space version) Let  $\mathcal{X} = [0, 1]$ . Define the transition kernel  $P(x, \cdot)$  as follows. If  $x = 1/m$  for some positive integer  $m$ , then  $P(x, \cdot) = x^2 \text{Uniform}[0, 1] + (1 - x^2) \delta_{1/(m+1)}(\cdot)$ . For all other  $x$ ,  $P(x, \cdot) = \text{Uniform}[0, 1]$ . Then the chain has stationary distribution  $\pi(\cdot) = \text{Uniform}[0, 1]$ , and it is  $\phi$ -irreducible (with respect to  $\pi$ ) and aperiodic. On the other hand, if  $X_0 = 1/m$  for some  $m \geq 2$ , then  $\mathbf{P}[X_n = 1/(m+n) \text{ for all } n] = \prod_{j=m}^{\infty} (1 - (1/j^2)) > 0$ , so that  $\|P^n(x, \cdot) - \pi(\cdot)\| \not\rightarrow 0$ . Hence, convergence fails to hold from the set  $G^C = \{1/2, 1/3, 1/4, \dots\}$ .

To rectify the problems of the null set  $G^C$ , we consider Harris recurrence, a concept developed by Harris (1956) and introduced to statisticians in the important works of Tierney (1994) and Chan and Geyer (1994) (see also Geyer, 1996).

**Definition 5.** A  $\phi$ -irreducible Markov chain with stationary distribution  $\pi(\cdot)$  is *Harris recurrent* if for all  $A \subseteq \mathcal{X}$  with  $\pi(A) > 0$ , and all  $x \in \mathcal{X}$ , we have  $\mathbf{P}(\tau_A < \infty \mid X_0 = x) = 1$ .

We prove the following equivalences.

**Theorem 6.** For a  $\phi$ -irreducible Markov chain with stationary probability distribution  $\pi(\cdot)$ , and period  $D \geq 1$ , the following are equivalent:

- (i) The chain is Harris recurrent.
- (ii) For all  $A \subseteq \mathcal{X}$  with  $\pi(A) > 0$ , and all  $x \in \mathcal{X}$ , we have  $\mathbf{P}(X_n \in A \text{ i.o.} \mid X_0 = x) = 1$ .  
[Here *i.o.* means “infinitely often”, i.e. for infinitely many different times  $n$ .]
- (iii) For all  $x \in \mathcal{X}$ ,  $\lim_{n \rightarrow \infty} \|(1/D) \sum_{r=1}^D P^{nD+r}(x, \cdot) - \pi(\cdot)\| = 0$ .
- (iv) For all  $x \in \mathcal{X}$ ,  $\mathbf{P}[\tau_G < \infty \mid X_0 = x] = 1$ , where  $G$  is as in Proposition 2.
- (v) For all  $x \in \mathcal{X}$ , and all  $A \in \mathcal{F}$  with  $\pi(A) = 1$ ,  $\mathbf{P}[\tau_A < \infty \mid X_0 = x] = 1$ .
- (vi) For all  $x \in \mathcal{X}$ , and all  $A \in \mathcal{F}$  with  $\pi(A) = 0$ ,  $\mathbf{P}[X_n \in A \text{ for all } n \mid X_0 = x] = 0$ .

**Proof.** (ii)  $\implies$  (i); (i)  $\implies$  (v); (v)  $\implies$  (iv); and (v)  $\iff$  (vi): Immediate.

(i)  $\implies$  (ii): Suppose to the contrary that (ii) does not hold. Then there is some  $A \subseteq \mathcal{X}$  with  $\pi(A) > 0$ , and  $x \in \mathcal{X}$  and  $N \in \mathbf{N}$ , such that  $\mathbf{P}(X_n \notin A \forall n \geq N \mid X_0 = x) > 0$ . Integrating over choices of  $y = X_N$ , this implies there is some  $y \in \mathcal{X}$  with  $\mathbf{P}(\tau_A = \infty \mid X_0 = y) > 0$ , contradicting (i).

(iv)  $\implies$  (iii): From Proposition 2, once the chain reaches  $G$ , it will then converge. The convergence in (iii) then follows. More formally, conditional on the first hitting time  $\tau_G$  and the corresponding chain value  $X_{\tau_G}$ , the chain will converge in total variation distance as in (iii). The conclusion (iii) then follows by integrating over all choices of  $\tau_G$  and  $X_{\tau_G}$ , and using the triangle inequality for total variation distance.

(iii)  $\implies$  (i): If  $\phi(A) > 0$  (where  $\phi(\cdot)$  is an irreducibility measure), then we must have  $\pi(A) > 0$  (see e.g. Lemma 3 of Rosenthal, 2001), so we have by (iii) that for all  $x \in \mathcal{X}$ ,  $\sum_{r=1}^D P^{nD+r}(x, A) \rightarrow D \pi(A) > 0$ , and in particular  $\sum_{n=1}^{\infty} P^n(x, A) = \infty$ . It then follows from Theorem 9.0.1 of Meyn and Tweedie (1993), using their definition of recurrence on p. 182, that we can find an absorbing subset  $H \subseteq \mathcal{X}$  such that the chain restricted to  $H$  is Harris recurrent, and  $\pi(H) = 1$ . Then  $(1/D) \sum_{r=1}^D P^{nD+r}(x, H) \rightarrow 1$ , so  $P^n(x, H) \rightarrow 1$ .

Hence, the chain will eventually reach  $H$  with probability 1. Since the chain restricted to  $H$  is Harris recurrent, it will then eventually reach any  $A$  with  $\pi(A) = 1$ , with probability 1, thus establishing (i). ■

For completeness, we note another method of verifying Harris recurrence (though we do not use it here). Given a Markov chain with stationary distribution  $\pi(\cdot)$ , a subset  $C \in \mathcal{F}$  is *small* if  $\pi(C) > 0$ , and there is an  $\epsilon > 0$  and a probability measure  $\nu(\cdot)$  on  $(\mathcal{X}, \mathcal{F})$ , such that  $P(x, A) \geq \epsilon \nu(A)$  for all  $A \in \mathcal{F}$  and  $x \in C$ . It follows easily that we must have  $\nu \ll \pi$ .

**Proposition 7.** (Meyn and Tweedie, 1993) *If a Markov chain with stationary distribution  $\pi(\cdot)$  has a small set  $C$  with the property that  $\mathbf{P}[X_n \in C \text{ i.o.} | X_0 = x] = 1$  for all  $x \in \mathcal{X}$ , then the chain is Harris recurrent.*

**Proof.** Using the splitting technique (e.g. Nummelin, 1984), we can regard the chain as proceeding by, each time it is in  $C$ , moving according to  $\nu(\cdot)$  with probability  $\epsilon$ . If the chain returns to  $C$  infinitely often, then with probability 1, it will eventually move according to  $\nu(\cdot)$ . Since  $\nu \ll \pi$ , this means it will eventually leave any set of null  $\pi$ -measure. Hence, the result follows from Theorem 6(vi). ■

Various drift conditions can be used to establish that  $\mathbf{P}[X_n \in C \text{ i.o.} | X_0 = x] = 1$  for all  $x \in \mathcal{X}$ , and thus establish Harris recurrence. For example, it follows from Meyn and Tweedie (1993, Theorem 13.0.1) that for  $\phi$ -irreducible chains, it suffices that there exists a measurable function  $V : \mathcal{X} \rightarrow (0, \infty)$  such that  $\mathbf{E}[V(X_1) | X_0 = x] \leq V(x) - 1 + b \mathbf{1}_C(x)$  for some  $b < \infty$ . Alternatively, it follows from Meyn and Tweedie (1993, Theorem 8.4.3) (see also Geyer, 1996) that for  $\phi$ -irreducible chains, Harris recurrence follows if there exists a measurable function  $V : \mathcal{X} \rightarrow (0, \infty)$  such that  $V^{-1}((0, \alpha])$  is small for all  $\alpha > 0$ , and such that  $\mathbf{E}[V(X_1) | X_0 = x] \leq V(x)$  for all  $x \in \mathcal{X} \setminus C$ .

**Remark.** We note that the null sets related to Harris recurrence are of an “extreme” kind, in the sense that the chain may fail entirely to converge from the null set. Less radically, one could consider chains which converge from everywhere, but which have a slower qualitative *rate* of convergence from some null set. For example, it should be possible to

construct Markov chains which are Harris recurrent, and which are geometrically ergodic, but converge at a sub-geometric rate from a certain null set of initial states. Or, chains which are polynomially ergodic at a particular polynomial rate  $\alpha$ , but fail to converge at the polynomial rate  $\alpha$  from some null set. Or, chains which are geometrically ergodic, but fail to converge from one null set, converge polynomially from another null set, converge sub-polynomially from a third null set, etc. In this context, Harris recurrence can be seen as one in a series of properties ensuring that “things aren’t worse when starting from a null set than when starting anywhere else”.

### 3. Full-dimensional Metropolis-Hastings Algorithms.

Let  $\mathcal{X}$  be some state space, with  $\sigma$ -algebra  $\mathcal{F}$ . Let  $\pi(\cdot)$  be a probability distribution on  $(\mathcal{X}, \mathcal{F})$  having unnormalised density function  $f : \mathcal{X} \rightarrow (0, \infty)$  with respect to some reference measure  $\nu(\cdot)$ , so that  $\int_{\mathcal{X}} f(x) \nu(dx) < \infty$  and

$$\pi(A) = \frac{\int_A f(x) \nu(dx)}{\int_{\mathcal{X}} f(x) \nu(dx)}, \quad A \in \mathcal{F}.$$

Note that we assume  $f > 0$  on  $\mathcal{X}$ , or equivalently that  $\mathcal{X}$  is defined to be the support of  $f$ . To avoid trivialities, we assume that  $\pi(\cdot)$  is not concentrated at a single state, i.e. that  $\pi\{x\} < 1$  for all  $x \in \mathcal{X}$ .

Let  $q : \mathcal{X} \times \mathcal{X} \rightarrow [0, \infty)$  be any jointly measurable function, such that  $\int q(x, y) \nu(dy) = 1$  for all  $x \in \mathcal{X}$ . Define the Markov kernel  $Q(x, \cdot)$  by  $Q(x, A) = \int_A q(x, y) \nu(dy)$  for  $x \in \mathcal{X}$ , and let

$$\alpha(x, y) = \min \left[ 1, \frac{f(y) q(y, x)}{f(x) q(x, y)} \right], \quad x, y \in \mathcal{X}$$

(with  $\alpha(x, y) = 1$  if  $f(x) q(x, y) = 0$ ).

The *full-dimensional Metropolis-Hastings algorithm* (Metropolis et al., 1953; Hastings, 1970; Tierney, 1994) proceeds as follows. Given that the chain is in the state  $X_n$  at time  $n$ , it generates a “proposal state”  $Y_{n+1} \sim Q(X_n, \cdot)$ . Then, with probability  $\alpha(X_n, Y_{n+1})$  it “accepts” this proposal and sets  $X_{n+1} = Y_{n+1}$ ; otherwise, with probability  $1 - \alpha(X_n, Y_{n+1})$ , it “rejects” this proposal and sets  $X_{n+1} = X_n$ . It is easy to check that  $\pi(\cdot)$  is then stationary for the Markov chain  $\{X_n\}$ .

Clearly, any such Markov chain can be decomposed as

$$P(x, A) = (1 - r(x)) M(x, A) + r(x) \delta_x(A), \quad x \in \mathcal{X}, \quad A \subseteq \mathcal{X},$$

where  $\delta_x(\cdot)$  is a point-mass at  $x$ ,  $r(x) = \int q(x, y)[1 - \alpha(x, y)]\nu(dy)$  is the probability of rejecting when starting at  $X_n = x$ , and  $M(x, \cdot)$  is the kernel conditional on moving (i.e., on  $X_{n+1} \neq X_n$ ). In particular, the probability distribution  $M(x, \cdot)$  is absolutely continuous with respect to  $\nu(\cdot)$ , for all  $x \in \mathcal{X}$ .

Regarding Harris recurrence, we have the following result, which was originally proved by Tierney (1994) using the theory of harmonic functions.

**Theorem 8.** (Tierney, 1994) *Every  $\phi$ -irreducible full dimensional Metropolis-Hastings algorithm is Harris recurrent.*

**Proof.** Since the chain is  $\phi$ -irreducible, and  $\pi\{x\} < 1$ , we must have  $r(x) < 1$  for all  $x \in \mathcal{X}$ . Suppose  $\pi(A) = 1$ . Then  $\pi(A^C) = 0$ , so since we are assuming that  $f > 0$  throughout  $\mathcal{X}$ , also  $\nu(A^C) = 0$ . Hence, by absolute continuity,  $M(x, A^C) = 0$ , i.e.  $M(x, A) = 1$ . It follows that, if the chain is at  $x$ , then it will eventually move according to  $M(x, \cdot)$ , at which point it will necessarily move into  $A$ . The result then follows from Theorem 6(v). ■

## 4. Metropolis-Within-Gibbs Algorithms.

We now define Metropolis-within-Gibbs Markov chains (Metropolis et al., 1953).

For simplicity, let  $\mathcal{X}$  be an open subset of  $\mathbf{R}^d$ , with Borel  $\sigma$ -algebra  $\mathcal{F}$ , and (unnormalised) target density  $f : \mathcal{X} \rightarrow (0, \infty)$  with  $\int_{\mathcal{X}} f(x) \lambda(dx) < \infty$  (where  $\lambda(\cdot)$  is  $d$ -dimensional Lebesgue measure). For  $1 \leq i \leq d$ , let  $q_i : \mathcal{X} \times \mathbf{R} \rightarrow [0, \infty)$  be jointly measurable, with  $\int_{-\infty}^{\infty} q_i(x, z) dz = 1$  for all  $x \in \mathcal{X}$  (where  $dz$  is one-dimensional Lebesgue measure).

Let  $Q_i(x, \cdot)$  be the Markov kernel on  $\mathbf{R}^d$  which replaces the  $i^{\text{th}}$  coordinate by a draw from the density  $q_i(x, \cdot)$ , while leaving the other coordinates unchanged. That is,

$$Q_i(x, S_{i,a,b}) = \int_a^b q_i(x, z) dz,$$

where

$$S_{i,a,b} = \{y \in \mathcal{X} : y_j = x_j \text{ for } j \neq i, \text{ and } a \leq y_i \leq b\}.$$

To avoid technicalities and special cases, assume that  $Q_i(x, \mathcal{X}) > 0$  for all  $x \in \mathcal{X}$  and  $1 \leq i \leq d$ , and also that each  $q_i$  is *symmetrically positive* in the sense that

$$q_i((x_1, \dots, x_{i-1}, y, x_{i+1}, \dots, x_d), z) > 0 \iff q_i((x_1, \dots, x_{i-1}, z, x_{i+1}, \dots, x_d), y) > 0.$$

For  $x, y \in \mathbf{R}^d$  and  $1 \leq i \leq d$ , let

$$\alpha_i(x, y) = \min \left[ 1, \frac{f(y) q_i(y, x)}{f(x) q_i(x, y)} \right]$$

(with  $\alpha_i(x, y) = 1$  if  $f(x) q_i(x, y) = 0$ ). Let  $P_i$  be the kernel which proceeds as follows. Given  $X_n$ , it generates a proposal  $Y_{n+1} \sim Q_i(X_n, \cdot)$ . Then, with probability  $\alpha_i(X_n, Y_{n+1})$  it accepts this proposal and sets  $X_{n+1} = Y_{n+1}$ ; otherwise, with probability  $1 - \alpha_i(X_n, Y_{n+1})$ , it rejects the proposal and sets  $X_{n+1} = X_n$ .

In terms of these definitions, the Metropolis-within-Gibbs Markov chain proceeds as follows. Random variables  $I_1, I_2, \dots$  taking values in  $\{1, 2, \dots, d\}$  are chosen according to some scheme. (The two most common schemes are random-scan, where  $\{I_n\}$  are i.i.d. uniform on  $\{1, 2, \dots, d\}$ , and deterministic-scan, where  $I_1 = 1, I_2 = 2, \dots, I_d = d, I_{d+1} = 1, \dots$ ) Then for  $n = 0, 1, 2, \dots$ , given  $X_n$ , the chain generates  $X_{n+1} \sim P_{I_{n+1}}(X_n, \cdot)$ . It is straightforward to verify that this chain has stationary distribution  $\pi(\cdot)$  given by

$$\pi(A) = \frac{\int_A f(x) \lambda(dx)}{\int_{\mathcal{X}} f(x) \lambda(dx)}, \quad A \in \mathcal{F}.$$

The above description defines Metropolis-Within-Gibbs chains as we shall study them. We can now ask, under what conditions are such chains Harris recurrent? One might think that a result similar to Theorem 8 holds for Metropolis-Within-Gibbs algorithms, at least when the target and proposal densities are continuous. However, to our initial surprise, this is false:

**Example 9.** We present a Metropolis-Within-Gibbs algorithm on an open subset  $\mathcal{X} \subseteq \mathbf{R}^2$ , with stationary distribution  $\pi(\cdot)$ , with continuous target and proposal densities, which is  $\phi$ -irreducible and aperiodic, but which fails to converge in distribution to  $\pi(\cdot)$  from an uncountable number of different starting points (having total  $\pi$ -measure zero, of course).

Let  $\mathcal{X} = \{(x_1, x_2) \in \mathbf{R}^2 : x_1 > 1\}$ , and define the function  $f : \mathcal{X} \rightarrow (0, \infty)$  by  $f(x_1, x_2) = (e/2) \exp(x_1 - |x_2|e^{2x_1})$  (so that  $\int_{\mathcal{X}} f = 1$ ). Let  $Q_1(x, \cdot)$  and  $Q_2(x, \cdot)$  be symmetric unit normal proposals, so that  $q_i(x, z) = (2\pi)^{-1/2} \exp(-(z - x_i)^2/2)$  ( $i = 1, 2$ ). Then clearly  $f$ ,  $q_1$ , and  $q_2$  are positive continuous functions; it follows that the chain is  $\phi$ -irreducible where  $\psi$  is Lebesgue measure.

Consider the random-scan (say) Metropolis-within-Gibbs Markov chain corresponding to these choices. We shall prove that this chain is not Harris recurrent. Indeed, let  $S = \{(x_1, 0) : x_1 > 1\}$  be the part of the line  $\{x_2 = 0\}$  which lies in  $\mathcal{X}$ . We claim that if the chain starts at any initial state in  $S$ , then there is positive probability that it will drift off to  $x_1 \rightarrow \infty$  without ever updating  $x_2$ , i.e. without ever leaving  $S$ . Then since  $\pi(S) = 0$ , it follows that if  $X_0 \in S$ , then the chain will fail to converge to  $\pi(\cdot)$ .

To establish the claim, consider first a Markov chain  $\{W_n\}$  equivalent to just the first coordinate of  $X_n$ , under just the kernel  $P_1$  (i.e., which proposes moves only in the  $x_1$  direction), restricted to the state space  $S$ . Now, on  $S$ , the density  $f$  is proportional to  $e^{x_1}$ . It follows that for any  $\delta > 0$  and  $x_1 > 1$ ,  $\alpha_1((x_1, 0), (x_1 - \delta, 0)) \leq e^{-\delta}$ , while  $\alpha_1((x_1, 0), (x_1 + \delta, 0)) = 1$ . That is, proposals to increase  $x_1$  will all be accepted, while a positive fraction of the proposals to decrease  $x_1$  will be rejected. It follows that on  $S$ , the kernel  $P_1$  has positive drift. Hence, there is  $c > 0$  such that for all  $x_1 > 1$ ,

$$\mathbf{P}[W_n \geq cn \text{ for all sufficiently large } n \mid W_0 = x_1] > 0.$$

On the other hand, the density  $f(x_1, x_2)$  as a function of  $x_2$  alone (i.e., with  $x_1$  regarded as a constant) is proportional to  $\exp(-|x_2|e^{2x_1})$ . It follows that the probability of accepting a proposal in the  $x_2$  direction is equal to  $\mathbf{E}[\exp(-|Z|e^{2x_1})]$  where  $Z \sim N(0, 1)$ , which is less than

$$\int_{-\infty}^{\infty} \exp(-|z|e^{2x_1}) dz = 2e^{-2x_1}.$$

Now, since  $\sum_n 2e^{-2cn} < \infty$ , it follows from the Borel-Cantelli Lemma (e.g. Rosenthal, 2000, Theorem 3.4.2) that there is positive probability that all proposed moves in the  $x_2$  direction will all be rejected. That is, for any  $x_1 > 1$ ,

$$\mathbf{P}[X_n \in S \text{ for all } n \mid X_0 = (x_1, 0)] > 0,$$

thus proving the claim. (We shall see in Corollary 18 below that the “problem” with this example is that the one-dimensional integral of  $f$  over the line  $\{x_2 = 0\}$  is infinite.)

**Remark.** In the above example, it is also possible, if desired, to modify  $f$  to decrease to 0 near the boundary  $\{x_1 = 1\}$ , to make  $f$  be continuous throughout  $\mathbf{R}^2$  (not just on  $\mathcal{X}$ ), without affecting the result.

To proceed, decompose  $P_i$  as  $P_i(x, \cdot) = [1 - r(x)] M_i(x, \cdot) + r(x) \delta_x(\cdot)$ , so that  $M_i(x, S)$  is the kernel corresponding to moving (i.e., both proposing and accepting) in the  $i^{\text{th}}$  direction.

**Lemma 10.** *Let  $(i_1, i_2, \dots, i_n)$  be any sequence of coordinate directions. Assume that each of the  $d$  coordinate directions appears at least once in the sequence  $(i_1, i_2, \dots, i_n)$ . Then  $M_{i_1} M_{i_2} \dots M_{i_n}$  is an absolutely continuous kernel, i.e. if  $A \in \mathcal{F}$  with  $\lambda(A) = 0$ , then  $(M_{i_1} M_{i_2} \dots M_{i_n})(x, A) = 0$  for all  $x \in \mathcal{X}$ .*

**Proof.** We shall compute a density for  $(M_{i_1} M_{i_2} \dots M_{i_n})(x, \cdot)$ ; the result then follows since every distribution having a density is absolutely continuous. Let

$$J = \{m : 1 \leq m \leq n, i_j \neq i_m \text{ for } m < j \leq n\},$$

i.e.  $J$  is the set of “last times we move in direction  $i$ ” for each coordinate  $i$ . (Thus,  $|J| = d$ .) For  $1 \leq m \leq n$ , let  $S_m \subseteq \mathbf{R}$  be any Borel subset, so that  $S = S_1 \times \dots \times S_d$  is an arbitrary measurable rectangle in  $\mathbf{R}^d$ . Then define subsets  $R_m \subseteq \mathbf{R}$  for  $1 \leq m \leq n$ , by saying that  $R_m = S_{i_m}$  if  $m \in J$ , otherwise  $R_m = \mathbf{R}$ .

We then compute that

$$(M_{i_1} M_{i_2} \dots M_{i_n})(x, S) =$$

$$\int_{R_1} \int_{R_2} \cdots \int_{R_n} q_{i_1}(x_1, x_2) \alpha(x_1, x_2) q_{i_2}(x_2, x_3) \alpha(x_2, x_3) \cdots \\ \cdots q_{i_{n-1}}(x_{n-1}, x_n) \alpha(x_{n-1}, x_n) dx_1 dx_2 \cdots dx_n.$$

It follows that the density of  $(M_{i_1} M_{i_2} \cdots M_{i_n})(x, \cdot)$  is given by the above formula, but with the integration over all of the variables  $\{x_j; j \in J\}$  all omitted. Hence,  $(M_{i_1} M_{i_2} \cdots M_{i_n})(x, \cdot)$  has a density, and is thus absolutely continuous. ■

From the Law of Total Probability, we therefore obtain:

**Corollary 11.** *If  $A$  has Lebesgue measure 0, then  $\mathbf{P}[X_n \in A \mid X_0 = x_0] \leq \mathbf{P}[D_n]$ , where  $D_n$  is the event that by time  $n$  the chain has not yet moved in each coordinate direction.*

This allows us to prove:

**Theorem 12.** *Consider a  $\phi$ -irreducible Metropolis-within-Gibbs Markov chain. Suppose that from any initial state  $x$ , with probability 1 the chain will eventually move at least once in each coordinate direction. Then the chain is Harris recurrent.*

**Proof.** The hypothesis implies that for all  $x \in \mathcal{X}$ ,  $\lim_{n \rightarrow \infty} \mathbf{P}[D_n \mid X_0 = x] = 0$ . Now, let  $\pi(A) = 0$ . Then, since  $f > 0$  on  $\mathcal{X}$ , we must also have  $\lambda(A) = 0$ . Hence, by Corollary 11, we must have

$$\mathbf{P}[X_n \in A \forall n \mid X_0 = x] \leq \lim_{n \rightarrow \infty} \mathbf{P}[X_n \in A \mid X_0 = x] \leq \lim_{n \rightarrow \infty} \mathbf{P}[D_n \mid X_0 = x] = 0.$$

The result then follows from Theorem 6(vi). ■

The classical Gibbs sampler (see e.g. Gelfand and Smith, 1990) is a special case of Metropolis-within-Gibbs in which the proposal densities are chosen so that  $\alpha(x, y) \equiv 1$ , i.e. so that all proposed moves are accepted. Now, with either the deterministic-scan or systematic-scan Gibbs sampler variants, it is certainly true that with probability 1, moves are eventually proposed in all directions. So, since  $\alpha(x, y) \equiv 1$ , then it follows that with probability 1 the chain will eventually move in all directions as well. Hence, from Theorem 12, we immediately obtain:

**Corollary 13.** (Tierney, 1994) *Every  $\phi$ -irreducible deterministic- or random-scan Gibbs sampler is Harris recurrent.*

## 5. Sub-Chains of Metropolis-Within-Gibbs Algorithms.

We now consider the extent to which Harris recurrence of the full chain can be “inherited” from Harris recurrence of various sub-chains. For a subset  $I = \{i_1, \dots, i_r\} \subseteq \{1, \dots, n\}$ , let  $P^{[I]}$  be the Markov kernel corresponding to the original Metropolis-within-Gibbs chain, except conditional on never moving in any coordinate directions other than the coordinate directions  $i_1, \dots, i_r$ . Call the collection of kernels  $P^{[I]}$  where  $|I| = d - 1$  the “ $(d - 1)$ -dimensional sub-chains”. These sub-chains can fail to be  $\phi$ -irreducible:

**Example 14.** Suppose that

$$\mathcal{X} = \{(x_1, x_2) \in \mathbf{R}^2 : 16 < x_1^2 + x_2^2 < 25\}$$

(an annulus or “donut-shaped” state space), and that the proposal kernels  $Q_i(x, \cdot)$  simply replace  $x_i$  by a draw from the Uniform $[x_i - 1, x_i + 1]$  distribution. Then the full Metropolis-within-Gibbs chain is  $\phi$ -irreducible, but the one-dimensional sub-chain along the line  $\{x_2 = 0\}$  breaks up into two distinct non-communicating intervals,  $(-5, -4)$  and  $(4, 5)$ , and is therefore not  $\phi$ -irreducible.

Harris recurrence is often defined just for  $\phi$ -irreducible chains (e.g. Meyn and Tweedie, 1993). We generalise as follows. Call a chain *piecewise Harris* if the state space  $\mathcal{X}$  can be partitioned into a disjoint union  $\mathcal{X} = \bigcup_{\alpha \in \mathcal{S}} \mathcal{X}_\alpha$ , where each  $\mathcal{X}_\alpha$  is closed, and the chain restricted to each  $\mathcal{X}_\alpha$  is Harris recurrent. Of course, if the partition consists of just a single  $\mathcal{X}_\alpha$ , then the full chain is Harris recurrent. The following proposition says that the piecewise Harris property often suffices.

**Proposition 15.** *If a Markov chain is piecewise Harris and is also  $\phi$ -irreducible, then it is Harris recurrent.*

**Proof.** Let  $\mathcal{X}_\beta$  be any non-empty element of the partition from the definition of piecewise Harris, and let  $\mathcal{X}_* = \mathcal{X} \setminus \mathcal{X}_\beta$ . If  $\phi(\mathcal{X}_*) > 0$ , then by  $\phi$ -irreducibility, for each  $x \in \mathcal{X}$  there exists  $n = n(x)$  with  $P^n(x, \mathcal{X}_*) > 0$ . Since  $\mathcal{X}_\beta$  is closed, this implies that  $\mathcal{X}_\beta$  is empty, contradicting our assumption. Hence,  $\phi(\mathcal{X}_*) = 0$ . Since  $\phi$  is non-zero, we must have  $\phi(\mathcal{X}_\beta) > 0$ . It then follows similarly that  $\mathcal{X}_*$  is empty, i.e. that  $\mathcal{X}_\beta = \mathcal{X}$ . Thus, the partition contains just a single element, so the chain is Harris recurrent.  $\blacksquare$

In terms of the piecewise Harris property, we have:

**Theorem 16.** *Consider a random-scan Metropolis-Within-Gibbs chain, as above. Suppose all the  $(d-1)$ -dimensional sub-chains, in every  $(d-1)$ -dimensional coordinate hyperplane, are all piecewise Harris. Then the full chain is piecewise Harris. (In particular, by Proposition 15, if the full chain is  $\phi$ -irreducible, then the full chain is Harris recurrent.)*

**Proof.** Consider any fixed initial state  $x_0 = (x_{0,1}, \dots, x_{0,d})$ . By Theorem 12, it suffices to show that with probability 1, when starting  $X_0 = x_0$ , the chain will eventually move in each coordinate direction.

Suppose to the contrary that this is false, and there is positive probability that the chain never moves in some direction, say (for notational simplicity) in direction  $d$ . Let  $H = \{y \in \mathcal{X} : y_j = x_{0,j} \text{ for } j \neq d\}$  be the hyperplane corresponding to never moving in the  $d^{\text{th}}$  direction.

Let  $I_n$  be the direction of the proposed move at time  $n$  of the full chain, and let  $A_n = 1$  if this move is accepted, otherwise  $A_n = 0$ . Let

$$C_{m,r} = \{w \in H; \mathbf{P}[I_m = d, A_m = 1 \mid X_0 = w] \geq 1/r\}.$$

That is,  $C_{m,r}$  is the set of states in  $H$  which have probability  $\geq 1/r$  of changing the  $d^{\text{th}}$  coordinate,  $m$  steps later, when moving according to the sub-chain.

By assumption,  $Q_d(x, \mathcal{X}) > 0$  and  $f(x) > 0$  for all  $x \in \mathcal{X}$ . This implies that the chain has positive probability, starting from any  $x \in H$ , to eventually move in the  $d^{\text{th}}$  direction, i.e. to leave  $H$ . Hence,

$$\bigcup_{m,r=1}^{\infty} C_{m,r} = H. \tag{1}$$

(In fact, it suffices to consider just  $m = 1$ , but we do not use that fact here.)

Consider now the sub-chain  $P^{[1,2,\dots,d-1]}$ , restricted to the hyperplane  $H$ . Since this sub-chain is piecewise Harris, we must have  $x_0 \in H_0$  for some closed subset  $H_0 \subseteq H$  such that the sub-chain restricted to  $H_0$  is Harris recurrent with respect to some non-zero measure  $\psi_j(\cdot)$ . From (1), there must be some  $m, r \in \mathbf{N}$  with  $\psi_j(C_{m,r}) > 0$ . It then follows from Theorem 6(ii) that with probability 1,  $C_{m,r}$  is hit infinitely often by the sub-chain. In other words, conditional on the full chain never moving in the  $d^{\text{th}}$  direction, it will enter  $C_{m,r}$  infinitely often.

However, each time the full chain visits  $C_{m,r}$ , it has probability  $\geq 1/r$  of moving in the  $d^{\text{th}}$  direction,  $m$  steps later. It follows that with probability 1, the full chain will eventually jump in the  $d^{\text{th}}$  direction, and hence leave  $H$ . This contradicts our assumption that the chain has positive probability of never leaving  $H$ . ■

Unfortunately, Theorem 16 still requires that we verify Harris recurrence of various sub-chains, which may be difficult. However, if the sub-chains of all dimensions all have stationary distributions, then no Harris recurrence needs to be checked:

**Corollary 17.** *Consider a random-scan Metropolis-Within-Gibbs chain as above. Suppose all the  $r$ -dimensional sub-chains in every  $r$ -dimensional coordinate hyperplane each have stationary probability distribution, for all  $1 \leq r \leq d$ . Then the full chain is piecewise Harris (as are all the sub-chains).*

**Proof.** Let  $T_r$  be the statement that all the sub-chains of dimension  $\leq r$  are all piecewise Harris. Then  $T_1$  holds by Theorem 8. Furthermore, from Theorem 16, for any  $r < d$ , if  $T_r$  holds, then  $T_{r+1}$  must hold. Hence, the result follows by induction. ■

We then have the following.

**Corollary 18.** *Consider a random-scan Metropolis-within-Gibbs Markov chain. Suppose the target density  $f$  has the property that its  $r$ -dimensional integral has finite Lebesgue integral, over every  $r$ -dimensional coordinate hyperplane of  $\mathcal{X}$ , for all  $1 \leq r \leq d$ . Then the full chain is piecewise Harris (as are all the sub-chains).*

**Proof.** In this case,  $f$  is an (unnormalised) density for a stationary probability distribution of each sub-chain on each hyperplane. (Note that the Lebesgue integral of  $f$  over the hyperplane must be positive, since we are assuming that  $\mathcal{X}$  is open and that  $f > 0$  on  $\mathcal{X}$ .) Hence, the result follows from Corollary 17.  $\blacksquare$

In the counter-example of Example 9, the one-dimensional  $x_1$ -chain fails to have a stationary distribution along the line  $\{x_2 = 0\}$ , since the integral of  $f$  along the line  $\{x_2 = 0\}$  is infinite.

A similar result to Corollary 18 appears as Theorem 1 of Chan and Geyer, 1994, under the assumption that each sub-chain is  $\phi$ -irreducible (which, as we have seen in Example 14, can easily fail to hold):

**Corollary 19.** (*Chan and Geyer, 1994*) *Consider a random-scan Metropolis-within-Gibbs Markov chain. Suppose the target density  $f$  has the property that its  $r$ -dimensional integral has finite Lebesgue integral, over every  $r$ -dimensional coordinate hyperplane of  $\mathcal{X}$ , for all  $1 \leq r \leq d$ . If the full chain and all the sub-chains are all  $\phi$ -irreducible, then the full chain is Harris recurrent.*

## 6. Transdimensional MCMC Algorithms.

In certain statistical setups (e.g. autoregressive models), the number of parameters is not fixed in advance. This means that the state space of possible parameter values is a (disjoint) union of spaces of different dimensions. Exploring such state spaces through MCMC algorithms requires the introduction of transdimensional MCMC. Transdimensional MCMC algorithms first appear in Norman and Filinov (1969) and Preston (1977); their introduction into modern statistical practice (under the name “reversible jump”) is due to the influential paper of Green (1995) (see also Tierney, 1998).

Suppose that for each  $m \in \mathcal{M} \subseteq \mathbf{N}$ , where  $|\mathcal{M}| > 1$  and usually  $|\mathcal{M}| = \infty$ , we have a space  $\mathcal{X}_m$  of dimension  $d_m$ , i.e.  $\mathcal{X}_m$  is an open subset of  $\mathbf{R}^{d_m}$ . We combine these different spaces into a single state space  $\mathcal{X}$  by setting  $\mathcal{X} = \bigcup_{m=1}^{\infty} (\{m\} \times \mathcal{X}_m)$ . Furthermore, suppose that on each  $\mathcal{X}_m$  we have an unnormalised target density function  $f_m : \mathcal{X}_m \rightarrow (0, \infty)$ , with

$\int_{\mathcal{X}_m} f_m < \infty$ . We then combine that into a single probability distribution  $\pi(\cdot)$  on  $\mathcal{X}$ , by choosing some  $p : \mathcal{M} \rightarrow (0, 1)$  with  $\sum_{m \in \mathcal{M}} p(m) = 1$ , and then setting

$$\pi(m, A) = p(m) \frac{\int_A f_m(x) \lambda_m(dx)}{\int_{\mathcal{X}} f_m(x) \lambda_m(dx)} \quad (2)$$

and using linearity; in (2),  $\lambda_m(\cdot)$  is Lebesgue measure on  $\mathbf{R}^{d_m}$ .

Transdimensional chains can proceed in a variety of ways (Green, 1995; Brooks et al., 2003). We first consider a general class which we call *full-dimensional transdimensional MCMC*. Fix some  $0 < a < 1$ , and some irreducible kernel  $R(m, \cdot)$  on  $\mathcal{M}$  such that  $R(m, m') > 0$  if and only if  $R(m', m) > 0$ . Then, at each iteration, with probability  $a$  the chain proposes a “between-model move” of the form of replacing  $(m, x)$  by  $(m', x')$ , where  $m' \sim R(m, \cdot)$ , and where  $x' \in \mathcal{X}_{m'}$  is generated by some complicated dimension-matching scheme (Green, 1995); this proposal is then accepted or rejected according to the usual Metropolis-Hastings scheme, except that now the formula for  $\alpha[(m, x), (m', x')]$  is more complicated and involves a Jacobian of the transformation used to generate  $x'$ . Otherwise, with probability  $1 - a$ , the chain leaves  $m$  fixed but proposes a “within-model move”, i.e. to replace  $x$  by  $x' \in \mathcal{X}_m$ , using a full-dimensional Metropolis-Hastings proposal on  $\mathcal{X}_m$ .

What about Harris recurrence? We have the following:

**Theorem 20.** *Consider a full-dimensional transdimensional MCMC algorithm as above. Let  $D$  be the event that no within-model move is ever accepted. Suppose  $\mathbf{P}[D | X_0 = (m, x)] = 0$  for all  $(m, x) \in \mathcal{X}$ . Then the algorithm is Harris recurrent.*

**Proof.** The proof is very similar to that of Theorem 8. Since  $\mathbf{P}[D | X_0 = (m, x)] = 0$ , the chain must eventually accept a within-model move. But since the within-model proposal distributions are full-dimensional, the probability of remaining in any set of  $\pi$ -measure 0 after such a move is equal to 0. The result thus follows from Theorem 6(vi). ■

**Remark.** Theorem 20 remains true regardless of the details of how the between-model moves are implemented, provided only that they preserve the stationarity of  $\pi(\cdot)$ .

**Remark.** Even without verifying the hypothesis of Theorem 20, it is true that once a full-dimensional transdimensional MCMC algorithm makes at least one within-model move, then since the within-model moves are full-dimensional, with probability 1 the chain will move to the set  $G$  of Proposition 2 and hence will then converge. The issue in Theorem 20 is whether or not such a within-model move will eventually occur with probability 1.

Now, Theorem 20 allows for the possibility that, from a null set, the model numbers  $m_n$  might have positive probability of converging to  $+\infty$  without ever accepting a within-model move, and this seems quite plausible. On the other hand, if the  $\{m_n\}$  process is recurrent, the situation is less clear, due to the complicated details of the  $(m, x) \rightarrow (m', x')$  mapping corresponding to the between-model moves. Conditional on never accepting a within-model move, even if the chain returns to  $\mathcal{X}_1$  (say) infinitely often, it might potentially be at “worse and worse” points within  $\mathcal{X}_1$  each time it returned, and thus have smaller and smaller probability of accepting within-model moves. So, even a chain in which  $\{m_n\}$  is recurrent could conceivably fail to be Harris recurrent. We state this as an open problem:

**Open Problem #1.** *Does there exist a  $\phi$ -irreducible full-dimensional transdimensional MCMC algorithm, as above, for which  $\mathbf{P}[m_n = 1 \text{ i.o.} \mid X_0 = (m, x)] = 1$  for all  $m \in \mathcal{M}$  and  $x \in \mathcal{X}_m$ , which fails to be Harris recurrent?*

More generally, transdimensional MCMC might not be full-dimensional. That is, the within-model moves might themselves be of the form of Metropolis-within-Gibbs. To model this, we proceed as in Brooks et al. (2003). We replace  $\mathcal{X}_m$  by  $\tilde{\mathcal{X}}_m \equiv \mathcal{X}_m \times [0, 1] \times [0, 1] \times \dots$ , with stationary distribution  $\tilde{\pi}_m = \pi_m \times \text{Uniform}[0, 1] \times \text{Uniform}[0, 1] \times \dots$ . We then let  $h_{ij} : \tilde{\mathcal{X}}_i \rightarrow \tilde{\mathcal{X}}_j$  be deterministic functions defined whenever  $R(i, j) > 0$ , such that  $h_{ji} = (h_{ij})^{-1}$ . The between-model moves are specified by saying that when the algorithm proposes changing  $m$  to  $m'$ , it simultaneously proposes changing  $x$  to  $h_{mm'}(x)$ .

A special case is when each  $h_{ij}$  function is simply the identity function, which is plausible if  $\mathcal{X}_m = [0, 1]^{d_m}$  for each  $m \in \mathcal{M}$ . More generally, we consider *coordinate-preserving transdimensional MCMC* in which each  $h_{ij}$  can be decomposed as

$$h_{ij} = h_{ij}^{(1)} \times h_{ij}^{(2)} \times \dots \tag{3}$$

where each  $h_{ij}^{(\ell)} : \mathbf{R} \rightarrow \mathbf{R}$  and its inverse are differentiable functions acting solely on the  $\ell^{\text{th}}$  coordinate. That is, the between-model moves modify each coordinate separately. The algorithm then proceeds, given a current state  $X_n = (m, x)$ , as follows. First, it replaces the coordinates  $d_m + 1, d_m + 2, \dots$  by fresh i.i.d. draws from the Uniform $[0, 1]$  distribution. (Of course, in practice we only need to generate such Uniform $[0, 1]$  draws when they are required. But from a theoretical perspective, it is simplest to pretend they are updated at each iteration.) Then, with probability  $a$  it proposes a between-model move as above, otherwise with probability  $1 - a$  it chooses one of the coordinates  $1, 2, \dots, d_m$  uniformly at random, and does a Metropolis-within-Gibbs proposal for that coordinate only.

**Theorem 21.** *Consider a  $\phi$ -irreducible transdimensional MCMC chain, which is coordinate-preserving as in (3). Suppose that for each  $(m, x) \in \mathcal{X}$ , when the chain starts at  $X_0 = (m, x)$ , then with probability 1 it eventually accepts at least one move in each of the coordinate directions  $1, 2, \dots, d_m$ . Then the chain is Harris recurrent.*

**Proof.** The proof is analogous to that of Theorem 12. The only difference is that we do not require a move to be accepted in the coordinate directions  $d_m + 1, d_m + 2, \dots$ , nor in the direction corresponding to the model index  $m$ . To justify this, note that since  $\mathcal{M}$  is countable, with  $p(m) > 0$  for all  $m \in \mathcal{M}$ , every distribution on  $\mathcal{M}$  is absolutely continuous. Also, when starting from  $X_0 = (m, x)$ , coordinates  $d_m + 1, d_m + 2, \dots$  are drawn from an absolutely continuous (Uniform) distribution automatically. So, in the context of Theorem 12, this is equivalent to having already moved in the coordinate directions  $d_m + 1, d_m + 2, \dots$  and in the direction of  $\mathcal{M}$ . Then, just as in Theorem 12, the chain will eventually leave any set of zero stationary measure. The result thus follows from Theorem 6(vi). ■

This leads to the question of Harris recurrence for transdimensional chains which are *coordinate-mixing*, i.e. not coordinate-preserving. Unfortunately, this situation is more complicated due to lack of control over composition of the  $h_{ij}$  functions. For manageability, call a transdimensional chain *dimension-controlling* if  $h_{ij}$  is the identity on coordinate directions  $\ell > \max(d_i, d_j)$ , i.e. if  $h_{ij}$  doesn't mix in any more dimensions than are necessary

for dimension-matching. Now, it seems that coordinate-mixing in the  $h_{ij}$  should only help the chain to avoid null sets. However, the difficulty is that the between-model moves could e.g. “swap” the values of two coordinates, so that updating each coordinate position once could correspond to updating one value twice, and the other value not at all. Thus, the situation is unclear, and we state this as an open problem:

**Open Problem #2.** *Consider a  $\phi$ -irreducible, coordinate-mixing, dimension-controlling transdimensional MCMC chain, as above. Suppose that for each  $(m, x) \in \mathcal{X}$ , when the chain starts at  $X_0 = (m, x)$ , then with probability 1 it eventually accepts at least one move in each of the coordinate directions  $1, 2, \dots, d_m$ . Does this imply that the chain is Harris recurrent?*

A positive answer to this question would show that general transdimensional chains, like other Metropolis-within-Gibbs chains, are Harris recurrent provided they eventually move at least once in each coordinate direction with probability 1.

**Acknowledgements.** We are grateful to Charlie Geyer for peaking our interest in Harris recurrence, and to Galin Jones for a helpful conversation. We thank the two anonymous referees for very detailed and insightful reports.

## REFERENCES

- P. Billingsley (1995), Probability and Measure, 3<sup>rd</sup> ed. John Wiley & Sons, New York.
- S.P. Brooks, P. Guidici, and G.O. Roberts (2003), Efficient construction of reversible jump Markov chain Monte Carlo proposal distributions (with discussion). J. Royal Stat. Soc. Series B **65**, 3–55.
- K.S. Chan and C.J. Geyer (1994), Discussion to Tierney (1994). Ann. Stat. **22**, 1747–1758.
- C.J. Geyer (1996), Harris Recurrence web page. Available at  
<http://www.stat.umn.edu/PAPERS/htmlprints/points/node39.html>
- P.J. Green (1995), Reversible jump MCMC computation and Bayesian model determination. Biometrika **82**, 711–732.

T. E. Harris (1956), The existence of stationary measures for certain Markov processes. In *Proceedings of the 3rd Berkeley Symposium on Mathematical Statistics and Probability, volume 2*, 113–124. University of California Press.

W.K. Hastings (1970), Monte Carlo sampling methods using Markov chains and their applications. *Biometrika* **57**, 97–109.

P.G. Hoel, S.C. Port, and C.J. Stone (1972), *Introduction to Stochastic Processes*. Waveland Press, Prospect Heights, Illinois.

G.L. Jones and J.P. Hobert (2001), Honest exploration of intractable probability distributions via Markov chain Monte Carlo. *Statistical Science* **16**, 312–334.

N. Metropolis, A. Rosenbluth, M. Rosenbluth, A. Teller, and E. Teller (1953), Equations of state calculations by fast computing machines. *J. Chem. Phys.* **21**, 1087–1091.

S.P. Meyn and R.L. Tweedie (1993), *Markov chains and stochastic stability*. Springer-Verlag, London.

G.E. Norman and V.S. Filinov (1969), Investigations of Phase Transitions by a Monte Carlo Method. *High Temperature* **7**, 216–222.

E. Nummelin (1984), *General irreducible Markov chains and non-negative operators*. Cambridge University Press.

C.J. Preston (1977), Spatial birth-death processes. *Bull. Int. Stat. Inst.* **46**, 371–391.

G.O. Roberts, J.S. Rosenthal, and P.O. Schwartz (1998), Convergence properties of perturbed Markov chains. *J. Appl. Prob.* **35**, 1–11.

G.O. Roberts and R.L. Tweedie (1999), Bounds on regeneration times and convergence rates for Markov chains. *Stoch. Proc. Appl.* **80**, 211–229. Correction: *Stoch. Proc. Appl.* **91** (2001), 337–338.

J.S. Rosenthal (1995a), Minorization Conditions and Convergence Rates for Markov Chain Monte Carlo. *J. Amer. Stat. Assoc.* **90**, 558–566.

J.S. Rosenthal (1995b), Convergence rates of Markov chains. *SIAM Review* **37**, 387–405.

J.S. Rosenthal (2000), *A first look at rigorous probability theory*. World Scientific Publishing Company, Singapore.

J.S. Rosenthal (2001), A review of asymptotic convergence for general state space

Markov chains. *Far East J. Th. Stat.* **5**, 37–50.

J.S. Rosenthal (2002), Quantitative convergence rates of Markov chains: A simple account. *Elec. Comm. Prob.* **7**, 123–128.

L. Tierney (1994), Markov chains for exploring posterior distributions (with discussion). *Ann. Stat.* **22**, 1701-1762.

L. Tierney (1998), A note on Metropolis-Hastings kernels for general state spaces. *Ann. Appl. Prob.* **8**, 1–9.