

Factorial ANOVA

More than one categorical
explanatory variable

This slide show is a free open source document.
See the last slide for copyright information.

Factorial ANOVA

- Categorical explanatory variables are called **factors**
- More than one at a time
- Designed for true experiments, but also useful with observational data
- If there are observations at all combinations of explanatory variable values, it's called a *complete* factorial design (as opposed to a fractional factorial).

The potato study

- Cases are potatoes
- Inoculate with bacteria, store for a fixed time period.
- Response variable is diameter of rotten spot in millimeters
- Two explanatory variables, randomly assigned
 - Bacteria Type (1, 2, 3)
 - Temperature (1=Cool, 2=Warm)

Two-factor design

	Bacteria Type		
Temp	1	2	3
1=Cool			
2=Warm			

Six treatment conditions

Factorial experiments

- Allow more than one factor to be investigated in the same study: Efficiency!
- Allow the scientist to see whether the effect of an explanatory variable *depends* on the value of another explanatory variable: Interactions
- Thank you again, Mr. Fisher.


Model: Data are normal with equal variance and conditional (cell) means $\mu_{i,j}$

	Bacteria Type			
Temp	1	2	3	
1=Cool	$\mu_{1,1}$	$\mu_{1,2}$	$\mu_{1,3}$	$\frac{\mu_{1,1} + \mu_{1,2} + \mu_{1,3}}{3}$
2=Warm	$\mu_{2,1}$	$\mu_{2,2}$	$\mu_{2,3}$	$\frac{\mu_{2,1} + \mu_{2,2} + \mu_{2,3}}{3}$
	$\frac{\mu_{1,1} + \mu_{2,1}}{2}$	$\frac{\mu_{1,2} + \mu_{2,2}}{2}$	$\frac{\mu_{1,3} + \mu_{2,3}}{2}$	μ



Tests

- Main effects: Differences among marginal means
- Interactions: Differences between differences (What is the effect of Factor A? **It depends** on level of Factor B.)


To understand the interaction, plot the means




Either Way




Non-parallel profiles = Interaction




Main effects for both variables, no interaction




Main effect for Bacteria only




Main Effect for Temperature Only



Both Main Effects, and the Interaction



Should you interpret the main effects?



Testing Contrasts

	Bacteria Type			
Temp	1	2	3	
1=Cool	$\mu_{1,1}$	$\mu_{1,2}$	$\mu_{1,3}$	$\frac{\mu_{1,1} + \mu_{1,2} + \mu_{1,3}}{3}$
2=Warm	$\mu_{2,1}$	$\mu_{2,2}$	$\mu_{2,3}$	$\frac{\mu_{2,1} + \mu_{2,2} + \mu_{2,3}}{3}$
	$\frac{\mu_{1,1} + \mu_{2,1}}{2}$	$\frac{\mu_{1,2} + \mu_{2,2}}{2}$	$\frac{\mu_{1,3} + \mu_{2,3}}{2}$	μ


- Differences between marginal means are definitely contrasts
- Interactions are also sets of contrasts

Interactions are sets of Contrasts

	Bacteria Type			
Temp	1	2	3	
1=Cool	$\mu_{1,1}$	$\mu_{1,2}$	$\mu_{1,3}$	$\frac{\mu_{1,1} + \mu_{1,2} + \mu_{1,3}}{3}$
2=Warm	$\mu_{2,1}$	$\mu_{2,2}$	$\mu_{2,3}$	$\frac{\mu_{2,1} + \mu_{2,2} + \mu_{2,3}}{3}$
	$\frac{\mu_{1,1} + \mu_{2,1}}{2}$	$\frac{\mu_{1,2} + \mu_{2,2}}{2}$	$\frac{\mu_{1,3} + \mu_{2,3}}{2}$	μ

- $H_0 : \mu_{1,1} - \mu_{2,1} = \mu_{1,2} - \mu_{2,2} = \mu_{1,3} - \mu_{2,3}$
- $H_0 : \mu_{1,2} - \mu_{1,1} = \mu_{2,2} - \mu_{2,1}$ and
 $\mu_{1,3} - \mu_{1,2} = \mu_{2,3} - \mu_{2,2}$

Interactions are sets of Contrasts



- $H_0 : \mu_{1,1} - \mu_{2,1} = \mu_{1,2} - \mu_{2,2} = \mu_{1,3} - \mu_{2,3}$
- $H_0 : \mu_{1,2} - \mu_{1,1} = \mu_{2,2} - \mu_{2,1}$ and
 $\mu_{1,3} - \mu_{1,2} = \mu_{2,3} - \mu_{2,2}$

Equivalent statements

- The effect of A depends upon B
- The effect of B depends on A

$$H_0 : \mu_{1,1} - \mu_{2,1} = \mu_{1,2} - \mu_{2,2} = \mu_{1,3} - \mu_{2,3}$$

$$H_0 : \mu_{1,2} - \mu_{1,1} = \mu_{2,2} - \mu_{2,1} \text{ and}$$

$$\mu_{1,3} - \mu_{1,2} = \mu_{2,3} - \mu_{2,2}$$

Three factors: A, B and C

- There are three (sets of) main effects: One each for A, B, C
- There are three two-factor interactions
 - A by B (Averaging over C)
 - A by C (Averaging over B)
 - B by C (Averaging over A)
- There is one three-factor interaction: $A \times B \times C$

Meaning of the 3-factor interaction

- The form of the $A \times B$ interaction depends on the value of C
- The form of the $A \times C$ interaction depends on the value of B
- The form of the $B \times C$ interaction depends on the value of A
- These statements are equivalent. Use the one that is easiest to understand.

To graph a three-factor interaction

- Make a separate mean plot (showing a 2-factor interaction) for each value of the third variable.
- In the potato study, a graph for each oxygen level.

Four-factor design

- Four sets of main effects
- Six two-factor interactions
- Four three-factor interactions
- One four-factor interaction: The nature of the three-factor interaction depends on the value of the 4th factor
- There is an F test for each one
- And so on ...

As the number of factors increases

- The higher-way interactions get harder and harder to understand
- All the tests are still tests of sets of contrasts (differences between differences of differences ...)
- But it gets harder and harder to write down the contrasts
- Effect coding becomes easier

Effect coding

Bact	B₁	B₂
1	1	0
2	0	1
3	-1	-1

Temperature	T
1=Cool	1
2=Warm	-1

$$E(Y|\mathbf{X} = \mathbf{x}) = \beta_0 + \beta_1 B_1 + \beta_2 B_2 + \beta_3 T + \beta_4 B_1 T + \beta_5 B_2 T$$

Interaction effects correspond to products of dummy variables

$$E(Y|\mathbf{X} = \mathbf{x}) = \beta_0 + \beta_1 B_1 + \beta_2 B_2 + \beta_3 T + \beta_4 B_1 T + \beta_5 B_2 T$$

- The A x B interaction: Multiply each dummy variable for A by each dummy variable for B
- Use these products as additional explanatory variables in the multiple regression
- The A x B x C interaction: Multiply each dummy variable for C by each product term from the A x B interaction
- Test the sets of product terms simultaneously

Make a table

$$E(Y|\mathbf{X} = \mathbf{x}) = \beta_0 + \beta_1 B_1 + \beta_2 B_2 + \beta_3 T + \beta_4 B_1 T + \beta_5 B_2 T$$

Bact	Temp	B_1	B_2	T	$B_1 T$	$B_2 T$	$E(Y \mathbf{X} = \mathbf{x})$
1	1	1	0	1	1	0	$\beta_0 + \beta_1 + \beta_3 + \beta_4$
1	2	1	0	-1	-1	0	$\beta_0 + \beta_1 - \beta_3 - \beta_4$
2	1	0	1	1	0	1	$\beta_0 + \beta_2 + \beta_3 + \beta_5$
2	2	0	1	-1	0	-1	$\beta_0 + \beta_2 - \beta_3 - \beta_5$
3	1	-1	-1	1	-1	-1	$\beta_0 - \beta_1 - \beta_2 + \beta_3 - \beta_4 - \beta_5$
3	2	-1	-1	-1	1	1	$\beta_0 - \beta_1 - \beta_2 - \beta_3 + \beta_4 + \beta_5$

Cell and Marginal Means

	Bacteria Type			
Tmp	1	2	3	
1=C	$\beta_0 + \beta_1 + \beta_3 + \beta_4$	$\beta_0 + \beta_2 + \beta_3 + \beta_5$	$\beta_0 - \beta_1 - \beta_2$ $+ \beta_3 - \beta_4 - \beta_5$	β_0 $+ \beta_3$
2=W	$\beta_0 + \beta_1 - \beta_3 - \beta_4$	$\beta_0 + \beta_2 - \beta_3 - \beta_5$	$\beta_0 - \beta_1 - \beta_2$ $- \beta_3 + \beta_4 + \beta_5$	β_0 $- \beta_3$
	$\beta_0 + \beta_1$	$\beta_0 + \beta_2$	$\beta_0 - \beta_1 - \beta_2$	β_0

We see

- Intercept is the grand mean
- Regression coefficients for the dummy variables are deviations of the marginal means from the grand mean
- What about the interactions?

$$E(Y|\mathbf{X} = \mathbf{x}) = \beta_0 + \beta_1 B_1 + \beta_2 B_2 + \beta_3 T + \beta_4 B_1 T + \beta_5 B_2 T$$

A bit of algebra shows

$$\mu_{1,1} - \mu_{2,1} = \mu_{1,2} - \mu_{2,2} \text{ is equivalent to } \beta_4 = \beta_5$$

$$\mu_{1,2} - \mu_{2,2} = \mu_{1,3} - \mu_{2,3} \text{ is equivalent to } 2\beta_4 = -\beta_5$$

$$\text{So } \beta_4 = \beta_5 = 0$$

Factorial ANOVA with effect coding is pretty automatic

- You don't have to make a table unless asked
- It always works as you expect it will
- Significance tests are the same as testing sets of contrasts
- Covariates present no problem. Main effects and interactions have their usual meanings, “controlling” for the covariates.
- Could plot the least squares means

Again

- Intercept is the grand mean
- Regression coefficients for the dummy variables are deviations of the marginal means from the grand mean
- Test of main effect(s) is test of the dummy variables for a factor.
- Interaction effects are regression coefficients corresponding to products of dummy variables.

Balanced vs. Unbalanced Experimental Designs

- Balanced design: Cell sample sizes are proportional (usually equal)
- Explanatory variables have zero relationship to one another
- Numerator SS in ANOVA are independent.
- Everything is nice and simple
- Most experimental studies are designed this way.
- As soon as somebody drops a test tube, it's no longer true

Analysis of unbalanced data

- When explanatory variables are related, there is potential ambiguity.
- A is related to Y, B is related to Y, and A is related to B.
- Who gets credit for the portion of variation in Y that could be explained by either A or B?
- With a regression approach, whether you use contrasts or dummy variables (equivalent), the answer is **nobody**.
- Think of full, restricted models.
- Equivalently, general linear test.

Some software is designed for balanced data

- The special purpose formulas are much simpler.
- They were very useful *in the past*.
- Since most real data are at least a little unbalanced, these formulas are a recipe for trouble.
- Most textbook data are balanced, so they cannot tell you what your software is really doing.
- R's `anova` and `aov` functions are designed for balanced data, though `anova` applied to `lm` objects can give you what you want if you use it with care.
- SAS `proc glm` is much more convenient. SAS `proc anova` is for balanced data. Avoid it.

Type I and Type III Tests

- `proc glm` displays both by default.
- Type III is the regression approach we know and love.
- We will use Type III and ignore Type I.
- But just for the record ...

Tests based on Type III SS

- Type III Sums of squares are sequential.
- In order of the effects in the `model` statement.
- Numerator of F-ratio is $(SSR_F - SSR_R)/s$, where the “restricted” model has all preceding terms, and the “full” model has those and also the effect being tested.
 - First term is tested not controlling for anything.
 - Second term is tested controlling for the first.
 - Third term is tested controlling for the first two.
 - And so on.
- But the denominator is MSE from the model with *all* effects.

Type I vs. Type III

- Type I test controls for all preceding effects.
- Type III test controls for *all* other effects.
- Type I and Type III tests of the *last* effect in the model are identical.
- For balanced data, Type I and Type III tests of all effects are the same.
- I can't remember what the Type II tests are.

Copyright Information

This slide show was prepared by Jerry Brunner, Department of Statistical Sciences, University of Toronto. It is licensed under a Creative Commons Attribution - ShareAlike 3.0 Unported License. Use any part of it as you like and share the result freely. These Powerpoint slides are available from the course website:

<http://www.utstat.toronto.edu/~brunner/oldclass/441s20>