

Last week

- ▶ Nuisance parameters $f(y; \psi, \lambda)$, $\ell(\psi, \lambda)$

- ▶ posterior marginal density

$$\pi_m(\psi) \doteq \frac{c}{\sqrt{(2\pi)^q}} e^{\ell_p(\psi) - \ell_p(\hat{\psi})} |j_p(\hat{\psi})|^{1/2} \frac{\pi(\psi, \hat{\lambda}_\psi)}{\pi(\hat{\psi}, \hat{\lambda})} \frac{|j_{\lambda\lambda}(\hat{\psi}, \hat{\lambda})|^{1/2}}{|j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|^{1/2}}$$

- ▶ $\ell_p(\psi) = \ell(\psi, \hat{\lambda}_\psi)$

- ▶ exact conditional density

$$f(y; \psi, \lambda) = \exp\{\psi^T s_1 + \lambda^T s_2 - k(\psi, \lambda)\} h(y)$$

- ▶

$$f(s_1 \mid s_2; \psi) \doteq \frac{c}{\sqrt{(2\pi)^q}} e^{\ell_p(\psi) - \ell_p(\hat{\psi})} |j_p(\hat{\psi})|^{-1/2} \frac{|j_{\lambda\lambda}(\hat{\psi}, \hat{\lambda})|^{-1/2}}{|j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|^{-1/2}}$$

... last week

Tail area approximation:

$$p(\psi) \doteq \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right),$$

$$r = \pm \sqrt{[2\{\ell(\hat{\theta}) - \ell(\theta)\}]} \text{ or } \pm \sqrt{[2\{\ell_p(\hat{\psi}) - \ell_p(\psi)\}]}$$

$$Q = q_B = -\ell'(\theta) j^{-1/2}(\hat{\theta}) \frac{\pi(\hat{\theta})}{\pi(\theta)} \quad \text{or}$$

$$-\ell_p(\hat{\psi}) j_p^{-1/2}(\hat{\psi}) \frac{\pi(\hat{\psi}, \hat{\lambda})}{\pi(\hat{\psi}, \hat{\lambda}_\psi)} \rho(\psi, \hat{\psi})$$

$$Q = q = (\hat{\theta} - \theta) j^{1/2}(\hat{\theta}) \text{ or } (\hat{\psi} - \psi) j_p^{1/2}(\hat{\psi}) \rho^{-1}(\psi, \hat{\psi})$$

$$(\hat{\theta} - \theta) j^{1/2} \quad \ell'(\theta) j^{-1/2} \pm \sqrt{2\{\ell(\hat{\theta}) - \ell(\theta)\}}$$

... last week

Tail area approximation:


$$p(\psi) \doteq \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right),$$

$$r = \pm \sqrt{[2\{\ell(\hat{\theta}) - \ell(\theta)\}]} \text{ or } \pm \sqrt{[2\{\ell_p(\hat{\psi}) - \ell_p(\psi)\}]}$$

$$\begin{aligned} Q &= q_B = -\ell'(\theta) j^{-1/2}(\hat{\theta}) \frac{\pi(\hat{\theta})}{\pi(\theta)} \quad \text{or} \\ &\quad -\ell_p(\hat{\psi}) j_p^{-1/2}(\hat{\psi}) \frac{\pi(\hat{\psi}, \hat{\lambda})}{\pi(\psi, \hat{\lambda}_\psi)} \rho(\psi, \hat{\psi}) \end{aligned}$$

$$Q = q = (\hat{\theta} - \theta) j^{1/2}(\hat{\theta}) \text{ or } (\hat{\psi} - \psi) j_p^{1/2}(\hat{\psi}) \rho^{-1}(\psi, \hat{\psi})$$

$$\rho(\psi, \hat{\psi}) = \frac{|j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|^{1/2}}{|j_{\lambda\lambda}(\hat{\psi}, \hat{\lambda})|^{1/2}}$$



P.int. Figure 2.3: Inference for shape parameter ψ of gamma sample of size $n = 5$. Left: profile log likelihood ℓ_p (solid) and the log likelihood from the conditional density of u given v (heavy). Right: likelihood root $r(\psi)$ (solid), Wald pivot $t(\psi)$ (dashes), modified likelihood root $r^*(\psi)$ (heavy), and exact pivot overlying $r^*(\psi)$. The horizontal lines are at $0, \pm 1.96$.

$$P(\psi) \propto \lambda^\psi e^{-\lambda x} x^{\psi-1} / \Gamma(\psi)$$

This week

1. elimination of nuisance parameters and adjusted profile likelihood
2. approximate inference in location and transformation models
3. tail area approximations again
4. tangent exponential model
5. examples

Nuisance parameters

- ▶ $\underline{\pi_m(\psi)} \doteq \frac{c}{\sqrt{(2\pi)^q}} e^{\ell_p(\psi) - \ell_p(\hat{\psi})} j_p^{1/2}(\hat{\psi}) \frac{\pi(\psi, \hat{\lambda}_\psi)}{\pi(\hat{\psi}, \hat{\lambda})} \frac{|j_{\lambda\lambda}(\hat{\psi}, \hat{\lambda})|^{1/2}}{|j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|^{1/2}}$
- ▶ $= \frac{c}{\sqrt{(2\pi)^q}} e^{\ell_A(\psi) - \ell_A(\hat{\psi})} j_p^{1/2}(\hat{\psi}) \frac{\pi(\psi, \hat{\lambda}_\psi)}{\pi(\hat{\psi}, \hat{\lambda})} \left\{ \frac{j_p^{1/2}}{j_A^{1/2}} \right\} \simeq 1$
- ▶ $\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|$
- ▶ $= \ell(\psi, \hat{\lambda}_\psi) - \dots$

$$f(s_1 \mid s_2; \psi) \doteq \frac{c}{\sqrt{(2\pi)^q}} |j_p(\hat{\psi})|^{-1/2} e^{\ell_p(\psi) - \ell_p(\hat{\psi})} \left\{ \frac{|j_{\lambda\lambda}(\hat{\psi}, \hat{\lambda})|}{|j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|} \right\}^{-1/2}$$

$$= \frac{c}{\sqrt{(2\pi)^q}} |j_p(\hat{\psi})|^{-1/2} e^{\ell_A(\psi) - \ell_A(\hat{\psi})}$$

$$\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|$$

$j_A - \ell_A''(\psi)$

Adjusted profile log-likelihood Modified profile

- ▶ $\ell_A(\psi) = \ell_p(\psi) + A(\psi) = \ell(\psi, \hat{\lambda}_\psi) + A(\psi)$
- ▶ $A(\psi)$ assumed to be $O_p(1)$
- ▶ generic form is $A_{FR}(\psi) = +\frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)| - \log |\frac{d(\lambda)}{d\hat{\lambda}_\psi}|$

Fraser, 2003



- ▶ closely related $A_{BN}(\psi) = -\frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)| + |\log \frac{\partial \hat{\lambda}}{\partial \hat{\lambda}_\psi}|$

SM §12.4.1, BN 1983

$$\hat{\lambda} = g(\hat{\lambda}_\psi, \hat{\psi}, \psi, a)$$

- ▶ if $i_{\psi\lambda}(\theta) = 0$, then $\hat{\lambda}_\psi = \hat{\lambda} + O_p(n^{-1})$, suggesting we ignore last term
- ▶ if ψ is scalar, then in principle we can find a parametrization (ψ, λ) in which $i_{\psi\lambda}(\theta) = 0$

$$l_A^*(\psi) = l_p(\psi) - \underset{\text{SM §12.4.2}}{\frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|} \quad \lambda \perp \psi$$

Log-likelihoods

- ▶ marginal $\ell_m(\psi) = \log f_m(t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi) f_c(s | t; \psi, \lambda)$$

- ▶ conditional $\ell_c(\psi) = \log f_c(s | t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi, \lambda) f_c(s | t; \psi)$$

- ▶ adjusted $\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)| \quad \lambda \perp \psi$
- ▶ refinements $\ell_{FR}(\psi), \ell_{BN}(\psi), \dots$

Sartori, 2003; Nov12 Prob 2.2

Log-likelihoods

- ▶ marginal $\ell_m(\psi) = \log f_m(t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi) f_c(s | t; \psi, \lambda)$$

- ▶ conditional $\ell_c(\psi) = \log f_c(s | t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi, \lambda) f_c(s | t; \psi)$$

- ▶ adjusted $\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)| \quad \lambda \perp \psi$
- ▶ refinements $\ell_{FR}(\psi), \ell_{BN}(\psi), \dots$

Sartori, 2003; Nov12 Prob 2.2

Log-likelihoods

- ▶ marginal $\ell_m(\psi) = \log f_m(t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi) f_c(s | t; \psi, \lambda)$$

- ▶ conditional $\ell_c(\psi) = \log f_c(s | t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi, \lambda) f_c(s | t; \psi)$$

- ▶ adjusted $\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)| \quad \lambda \perp \psi$
- ▶ refinements $\ell_{FR}(\psi), \ell_{BN}(\psi), \dots$

Sartori, 2003; Nov12 Prob 2.2

Log-likelihoods

- marginal $\ell_m(\psi) = \log f_m(t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi) f_c(s | t; \psi, \lambda)$$

- conditional $\ell_c(\psi) = \log f_c(s | t; \psi)$

$$f(y; \psi, \lambda) \propto f_m(t; \psi, \lambda) f_c(s | t; \psi)$$

- adjusted $\ell_A(\psi) = \ell_p(\psi) - \frac{1}{2} \log |j_{\lambda\lambda}(\psi, \hat{\lambda}_\psi)|$ $\lambda \perp \psi$

$$\ell'_A(\hat{\lambda}_A) = O((\hat{\lambda}_A - \lambda)^{-1}) j''_{\lambda\lambda}(\hat{\lambda}_A) \sim N(0, 1)$$

- refinements $\ell_{FR}(\psi), \ell_{BN}(\psi), \dots$ $\{1 + O(n^{-1/2})\}$

$d = d_n$
MLE's
not conc.
or
not effic.

$$(\psi, \lambda)$$

 $x \in \mathbb{R}^{d-q}$

$$q=1 \quad \lambda \in \mathbb{R}^{d-1} \quad n \quad O(n^{-?})$$

Sartori, 2003; Nov12 Prob 2.2





Figure 12.11
Likelihood analysis of Danish fire data. Left: variation of $\hat{\sigma}_\xi$ (solid) and $\hat{\lambda}_\xi$ (dashes) as a function of the shape parameter ξ . The blobs show the maximum likelihood estimates. The dotted line is the profile log likelihood $\ell_p(\xi)$ and the vertical lines mark the limits of a 0.99 confidence interval for ξ . The orthogonal parameter $\hat{\lambda}_\xi$ varies much less than the non-orthogonal parameter $\hat{\sigma}_\xi$ over the range of ψ considered. Right: profile log likelihood (solid) and modified profile log likelihood (dashes) for 0.99 quantile ψ of generalized Pareto distribution. The horizontal line determines the limits of a 0.95 confidence interval for ψ .

Example 12.25 (Generalized Pareto distribution) The expected information matrix for an observation with distribution function (6.38) is

$$i^*(\xi, \sigma) = \frac{1}{\sigma^2(1 + \xi)(1 + 2\xi)} \begin{pmatrix} 2\sigma^2 & \sigma \\ \sigma & 1 + \xi \end{pmatrix}.$$


Hence a parameter $\lambda = \lambda(\xi, \sigma)$ orthogonal to the shape parameter ξ satisfies the partial differential equation

$$f(s | t; \psi)$$

544

N. SARTORI

$$\begin{aligned} l_c(\psi) \\ l_A(\psi) \\ l_P(\psi) \end{aligned}$$




$$(y_{1i}, y_{2i}) \sim \text{Bin}$$

$$y_{1i} \sim B(m, p_{1i})$$

$$y_{2i} \sim B(m, p_{2i})$$

$$i=1, \dots, 100$$

Fig. 1: Example 3. Inference about the common odds ratio in pairs of binomial observations. Relative loglikelihoods for ψ : profile (dotted), modified profile (dashed) and conditional (solid). The horizontal dashed line gives the 0.90 confidence interval. We consider $q = 100$, $m = 5$, $v_i = 3$ ($i = 1, \dots, q$), and three values of u .



$$\psi = \log \left(\frac{p_{1i}/(1-p_{1i})}{p_{2i}/(1-p_{2i})} \right)$$

$$\lambda_i = \log \left(\frac{p_{2i}}{1-p_{2i}} \right)$$

Marginal inference

- ▶ conditional inference
 - linear exponential families saddlepoint approx
 - ▶ marginal inference?
 - ▶ Bayesian posterior Laplace approx
 - ▶ transformation models
 - ▶ exponential families – non-linear parameter of interest
 - ▶ tangent exponential models
 - ▶ everything else

Location model

- ▶ location model $f(y_i; \theta) = f_0(y_i - \theta)$, $y_i, \theta \in \mathbb{R}$ $i = 1, \dots, n$
- ▶ $f(y; \theta) = \prod f_0(y_i - \theta)$
- ▶ $\ell(\theta; y) = \sum \log f_0(y_i - \theta) = \sum_{i=1}^n g_0(y_i - \theta)$
- ▶ change of variable $(y_1, \dots, y_n) \leftrightarrow (\hat{\theta}, \underline{a}_1, \dots, \underline{a}_n)$ $\underline{a} \in \mathbb{R}^{n-1}$
- ▶ $a_i = y_i - \hat{\theta}$
- ▶ $f(\hat{\theta}, \underline{a}; \theta) = \prod f_0(a_i + \hat{\theta} - \theta) | \underline{J} |$

$$\ell'(\hat{\theta}) = 0 \Rightarrow -\sum g'_0(y_i - \hat{\theta}) = 0 = \sum_{i=1}^n g'(a_i) = 0$$

... location models

► $f(\hat{\theta}, a; \theta) = \frac{\pi f_0(a_i + \hat{\theta} - \theta) |J|}{\int \pi f_0(a_i + \hat{\theta} - \theta) |J| d\hat{\theta}}$

► $f(a; \theta) = \frac{\pi f_0(a_i + \hat{\theta} - \theta) |J| d\hat{\theta}}{f(y) \propto f_c(\hat{\theta}|a; \theta) f_m(a)}$

► $f(\hat{\theta} | a; \theta) = \frac{\pi f_0(a_i + \hat{\theta} - \theta)}{\int \pi f_0(a_i + \hat{\theta}) dt}$

ntbc $= \dots = \frac{L(\theta; \hat{\theta}, \underline{a})}{\int L(\theta; \hat{\theta}, a) d\hat{\theta}}$

$|J| = h(\underline{a}) \left| \frac{\partial(y)}{\partial(\hat{\theta}; \underline{a})} \right|$

$$f(\hat{\theta} | \alpha) = \frac{L(\theta; \hat{\theta}, \alpha)}{\int L(\vartheta; \hat{\theta}, \alpha) d\vartheta}$$

$$\cancel{V_{\text{apfakt}}} \doteq c \cdot \underbrace{|j(\hat{\theta})|^{\frac{1}{2}}}_{l(\theta)} e^{\underbrace{l(\theta) - l(\hat{\theta})}_{\text{}}}$$

$$l(\theta; y) \\ l(\theta) = l(\theta; \hat{\theta}, \alpha)$$

$$= \sum \log f_0(a_i + \hat{\theta} - \theta_k)$$

Approximate p-values

$$= -\frac{1}{2} \eta^2$$

$$\begin{aligned}\int_{-\infty}^{\hat{\theta}} f(\hat{\vartheta} | a; \vartheta) d\hat{\vartheta} &= c \int_{-\infty}^{\hat{\theta}} |j(\hat{\vartheta})|^{1/2} e^{\ell(\vartheta; \hat{\vartheta}, a) - \ell(\hat{\vartheta}; \hat{\vartheta}, a)} d\hat{\vartheta} \\ &= \int_0^r c e^{-\frac{1}{2}r^2} |j(\hat{\vartheta})|^{1/2} \frac{dr}{\ell_{;\hat{\vartheta}}(\hat{\vartheta}; \hat{\vartheta}, a) - \ell_{;\hat{\vartheta}}(\vartheta; \hat{\vartheta}, a)} d\hat{\vartheta}\end{aligned}$$

$$\begin{aligned}-r dr \\ = \ell_{;\hat{\vartheta}}(\theta; \hat{\vartheta}, a) \\ = \ell_{;\hat{\vartheta}}(\hat{\theta}; \hat{\theta}, a)\end{aligned}$$

$$-\ell_{;\hat{\theta}}(\hat{\theta}; \hat{\theta}, a)$$

$$\begin{aligned}-\ell_{;\hat{\theta}}(\hat{\theta}; \hat{\theta}, a) &= \{\ell_{;\hat{\theta}}(\hat{\theta}; \hat{\theta}, a) - \ell_{;\hat{\theta}}(\theta; \hat{\theta}, a)\} j(\hat{\theta})^{-1/2} \\ &= \ell_\theta(\theta) j(\hat{\theta})^{-1/2}\end{aligned}$$

↑ Simplifies to →

location model

... approximate p-values

$$f(\hat{\theta} | a; \theta) = c \cdot j(\hat{\theta}) \cdot e^{q(\theta) - l(\hat{\theta})}$$

$$\Leftrightarrow (\hat{\theta}, a) \quad f_m(a; \theta) = f_m(a)$$

approx. anc. $\{1 + O(\bar{n})\}$

$$\Rightarrow F(\hat{\theta} | a; \theta) = \Phi(r^*)$$

$$= \Phi\left(r + \frac{1}{n} \log \frac{q}{r}\right)$$

$$q = \left\{ l_{,\hat{\theta}}(\theta; \hat{\theta}, a) - l_{,\hat{\theta}}(\hat{\theta}; \hat{\theta}, a) \right\}^{-1/2}$$

BUT

ancillary

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} | a; \sigma, \beta) f_m(a) \quad a =$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again


Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = \underbrace{f_c(\hat{\beta}, \hat{\sigma} | a; \sigma, \beta)}_{e^{-\frac{1}{2} \sum_j t_j^2 / v_j}} f_m(a) \quad a = \frac{(y_i - x_i^T \hat{\beta})}{\hat{\sigma}}$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} \mid a; \sigma, \beta) f_m(a) \quad a =$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} | a; \sigma, \beta) f_m(a) \quad a =$  
- ▶ parameter of interest β_j , say 
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} \mid a; \sigma, \beta) f_m(a) \quad a =$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i$, $\epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} | a; \sigma, \beta) f_m(a)$ $a = \hat{\beta}$ = mle
or $(x^T x)^{-1} x^T y$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$ either $j^{-1} (\hat{\beta})$ or $\frac{j^{-1} (\hat{\beta})}{\sigma^2 (x^T x)^{-1} \text{Var}_0(\epsilon)}$
- ▶ p -value functions using Laplace approximation again

Nuisance parameters

- ▶ Regression-scale models: $y_i = x_i^T \beta + \sigma \epsilon_i, \quad \epsilon_i \sim f_0(\cdot)$
- ▶ $f(y; \beta, \sigma^2) = f_c(\hat{\beta}, \hat{\sigma} | a; \sigma, \beta) f_m(a) \quad a =$
- ▶ parameter of interest $\hat{\beta}_j$, say
- ▶ $t_j = \frac{\hat{\beta}_j - \beta_j}{v_j}$ has marginal density free of $\beta_{(j)}, \sigma$
- ▶ $t_{p+1} = \log \hat{\sigma} - \log \sigma$ has marginal density free of β
- ▶ $v_j = \text{Var}(\hat{\beta}_j) =$
- ▶ p -value functions using Laplace approximation again

BDR Ch5 marg (marginal)

These are all special cases

- ▶ to compute p -value we need to integrate a density approximation
- ▶ this density approximation is to either a marginal or a conditional density
- ▶ the integration involves the derivative of the log-likelihood, with respect to the data
- ▶ simplify the density approximation to incorporate only this first derivative
- ▶ use the resulting simpler form as the basis for approximate likelihood based inference

These are all special cases

- ▶ to compute p -value we need to integrate a density approximation
- ▶ this density approximation is to either a marginal or a conditional density
- ▶ the integration involves the derivative of the log-likelihood, with respect to the data
- ▶ simplify the density approximation to incorporate only this first derivative
- ▶ use the resulting simpler form as the basis for approximate likelihood based inference

These are all special cases

- ▶ to compute p -value we need to integrate a density approximation
- ▶ this density approximation is to either a marginal or a conditional density
- ▶ the integration involves the derivative of the log-likelihood, with respect to the data
- ▶ simplify the density approximation to incorporate only this first derivative
- ▶ use the resulting simpler form as the basis for approximate likelihood based inference

These are all special cases

- ▶ to compute p -value we need to integrate a density approximation
- ▶ this density approximation is to either a marginal or a conditional density
- ▶ the integration involves the derivative of the log-likelihood, with respect to the data
- ▶ simplify the density approximation to incorporate only this first derivative
- ▶ use the resulting simpler form as the basis for approximate likelihood based inference

These are all special cases

- ▶ to compute p -value we need to integrate a density approximation
- ▶ this density approximation is to either a marginal or a conditional density
- ▶ the integration involves the derivative of the log-likelihood, with respect to the data
- ▶ simplify the density approximation to incorporate only this first derivative
- ▶ use the resulting simpler form as the basis for approximate likelihood based inference

Tangent exponential model $y \sim f(y; \theta) \leftarrow \theta \in \mathbb{R}^d$

y^0

► $f_{TEM}(s; \theta) = c e^{\ell\{\varphi(\theta); y^0\} + \varphi(\theta)^T s} |j_{\varphi\varphi}\{\hat{\varphi}(s); s\}|^{-1/2}$

$y \in \mathbb{R}^n$

► $p\text{-value} = \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right)$

$y \downarrow s \in \mathbb{R}^d$
 $\theta \rightarrow \varphi(\theta)$

► $Q = \{\varphi(\hat{\theta}) - \varphi(\theta)\} J_{\varphi\varphi}^{-1/2}(\hat{\varphi}) = \frac{\varphi(\hat{\theta}) - \varphi(\theta)}{\varphi_\theta(\hat{\theta})} J_{\theta\theta}^{1/2}(\hat{\theta})$

$\in \mathbb{R}^d$

► $Q = \frac{|\varphi(\hat{\theta}) - \varphi(\hat{\theta}_\psi)| |\varphi_\lambda(\hat{\theta}_\psi)|}{|\varphi_\theta(\hat{\theta})|} \frac{|j(\hat{\theta})|^{1/2}}{|j_{\lambda\lambda}(\hat{\theta}_\psi)|^{1/2}}$

Tangent exponential model

- ▶ $f_{TEM}(s; \theta) = c e^{\ell\{\varphi(\theta); y^0\} + \varphi(\theta)^T s} |j_{\varphi\varphi}\{\hat{\varphi}(s); s\}|^{-1/2}$
- ▶ $p\text{-value} = \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right)$
- ▶ $Q = \{\varphi(\hat{\theta}) - \varphi(\theta)\} J_{\varphi\varphi}^{-1/2}(\hat{\varphi}) = \frac{\varphi(\hat{\theta}) - \varphi(\theta)}{\varphi_\theta(\hat{\theta})} J_{\theta\theta}^{1/2}(\hat{\theta})$
- ▶ $Q = \frac{|\varphi(\hat{\theta}) - \varphi(\hat{\theta}_\psi)|}{|\varphi_\theta(\hat{\theta})|} \frac{|j(\hat{\theta})|^{1/2}}{|j_{\lambda\lambda}(\hat{\theta}_\psi)|^{1/2}}$

Tangent exponential model

- ▶ $f_{TEM}(s; \theta) = c e^{\ell\{\varphi(\theta); y^0\} + \varphi(\theta)^T s} |j_{\varphi\varphi}\{\hat{\varphi}(s); s\}|^{-1/2}$
- ▶ $p\text{-value} = \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right)$
- ▶ $Q = \{\varphi(\hat{\theta}) - \varphi(\theta)\} J_{\varphi\varphi}^{-1/2}(\hat{\varphi}) = \frac{\varphi(\hat{\theta}) - \varphi(\theta)}{\varphi_\theta(\hat{\theta})} J_{\theta\theta}^{1/2}(\hat{\theta}) \quad \theta \in \mathbb{R}$
no mits.
- ▶ $Q = \frac{|\varphi(\hat{\theta}) - \varphi(\hat{\theta}_\psi)| \varphi_\lambda(\hat{\theta}_\psi)|}{|\varphi_\theta(\hat{\theta})|} \frac{|j(\hat{\theta})|^{1/2}}{|j_{\lambda\lambda}(\hat{\theta}_\psi)|^{1/2}}$

Tangent exponential model

- ▶ $f_{TEM}(s; \theta) = c e^{\ell\{\varphi(\theta); y^0\} + \varphi(\theta)^T s} |j_{\varphi\varphi}\{\hat{\varphi}(s); s\}|^{-1/2}$
- ▶ $p\text{-value} = \Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{Q}{r}\right)$
- ▶ $Q = \{\varphi(\hat{\theta}) - \varphi(\theta)\} J_{\varphi\varphi}^{-1/2}(\hat{\varphi}) = \frac{\varphi(\hat{\theta}) - \varphi(\theta)}{\varphi_\theta(\hat{\theta})} J_{\theta\theta}^{1/2}(\hat{\theta})$
 $\approx |\ell(\hat{\theta}) - \ell(\hat{\theta}_\psi)| \quad \left(\frac{\ell(\hat{\theta}) - \ell(\hat{\theta}_\psi)}{J_{\theta\theta}(\hat{\theta}_\psi)} \right) |$ BN
not n
- ▶
$$Q = \frac{|\varphi(\hat{\theta}) - \varphi(\hat{\theta}_\psi)|}{|\varphi_\theta(\hat{\theta})|} \frac{|\varphi_\lambda(\hat{\theta}_\psi)|}{|j_{\lambda\lambda}(\hat{\theta}_\psi)|^{1/2}} \left| \begin{array}{c} \hat{\varphi}_1 - \tilde{\varphi}_1 \\ \vdots \\ \hat{\varphi}_d - \tilde{\varphi}_d \end{array} \right|^m$$

$$\Theta = (\psi, \lambda)$$

... tangent exponential model

$$f_{TEM}(s; \theta) = c e^{\ell\{\varphi(\theta); y^0\} + \varphi(\theta)^T s} |j_{\varphi\varphi}\{\hat{\varphi}(s); s\}|^{-1/2}$$

any 'smooth' model

$$F \leftarrow \Phi(n^*) = \Phi\left(n + \frac{1}{2} \log \frac{Q}{n}\right)$$

$$Q = Q \leftarrow \varphi(l, \varphi)$$

$$n = \sqrt{\ell_p(\hat{\psi}) - \ell_p(\psi)}$$

.. tangent exponential model

$$\varphi(\theta)^\top = \ell; v(\theta; y) = \sum_{i=1}^n \frac{\partial \ell(\theta; y)}{\partial y_i} \Big|_{y=y^*} V_i$$

$\in \mathbb{R}^d$ $\in \mathbb{R}^n$

$$i = 1, \dots, n$$

$$V_i = (\dots)_d$$

$$e^{\varphi'y - c(\varphi)}$$

$$h(y)$$

$$\triangleright V_i = - \left(\frac{\partial z_i}{\partial y_i} \right)^{-1} \frac{\partial z_i}{\partial \theta} \Big|_{(\hat{\theta}, y)}$$

BDR, Ch. 8.4, 8.5

—

.. tangent exponential model

$$\varphi(\theta) = \ell; v(\theta; y) = \sum_{i=1}^n \frac{\partial \ell(\theta; y)}{\partial y_i} V_i$$

$$v_i = - \left(\frac{\partial z_i}{\partial y_i} \right)^{-1} \frac{\partial z_i}{\partial \theta} \Bigg|_{(\hat{\theta}, y)}$$

← formula
for V

$z_i = z(y_i, \theta)$ whose dist is known
(prior)

BDR, Ch. 8.4, 8.5

Examples

- ▶ location-scale model: $y_i = \mu + \sigma\epsilon_i, z_i = \epsilon_i = (y_i - \mu)/\sigma$
- ▶ $V = -\left(\frac{\partial z_i}{\partial y_i}\right)^T \left(\frac{\partial z_i}{\partial \theta}\right)$ $\left| \begin{array}{l} \epsilon_i \sim f_0 \\ \text{Y, } \hat{\theta} \\ n \times 2 \end{array} \right.$ $\xrightarrow{\quad \quad \quad \quad \quad}$ $V = \begin{pmatrix} 1 & (y_1 - \hat{\mu})/\hat{\sigma} \\ 1 & (y_2 - \hat{\mu})/\hat{\sigma} \\ \vdots & \vdots \\ 1 & (y_n - \hat{\mu})/\hat{\sigma} \end{pmatrix}$
- ▶ location-regression model $y_i = x_i^T \beta + \sigma\epsilon_i, z_i =$
- ▶ $V =$

Example: top quark

- ▶ model $Y \sim \text{Poisson}(\mu + b)$, b known

- ▶ data $y = 27$ $b = 6.7$

- ▶ mid p -value

$$\underline{\Pr(Y > 27)} + \underline{\frac{1}{2}\Pr(Y = 27)} = \underline{18.45 \times 10^{-10}}$$

- ▶ approximation: $\Phi(r^*) = \Phi\left(r + \frac{1}{r} \log \frac{q}{r}\right) = \underline{15.85 \times 10^{-10}}$

{ continuity correction: $\Phi\left(r^*(y + \frac{1}{2})\right) = 32.36 \times 10^{-10}$

$$\Pr(Y \geq 27) = 29.83 \times 10^{-10}$$

Abe et al., 1995

... top quark

	$\text{pr}(Y \geq y^0; \mu = 0)$	$\text{pr}(Y > y^0; \mu = 0)$
Exact	29.83×10^{-10}	7.06×10^{-10}
Mid- p	18.45×10^{-10}	
$\Phi(r^*)$	15.85×10^{-10}	
$N(\theta, \theta)$	4.45×10^{-4}	2.21×10^{-5}
$N(\theta, \hat{\theta})$	1.02×10^{-4}	4.68×10^{-5}

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

Exact inference $\left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

Evans et al. 1999

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

Exact inference $\left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

Evans et al. 1999

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

Exact inference $\left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

Evans et al. 1999

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

$$\text{Exact inference } \left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

Evans et al. 1999

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

$$\text{Exact inference } \left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

Evans et al. 1999

Two-sample comparison

data on cost of treatment: standard vs new treatment

Group 1	30	172	210	212	335	489	651	1263
	1875	2213	2998	4935				
Group 2	121	172	201	214	228	261	278	279
	561	622	694	848	853	1086	1110	1243

model $Y_{1i} \sim \text{Exp}(\mu_1)$, $Y_{2i} \sim \text{Exp}(\mu_2)$ $\psi = \mu_1/\mu_2$

Exact inference $\left(\frac{\bar{Y}_{1\cdot}}{\mu_1} \right) / \left(\frac{\bar{Y}_{2\cdot}}{\mu_2} \right) \sim F_{2n, 2m}$

$E(\pi^*)$

$\pi^* = \pi + \dots$

exact 95% confidence interval for ψ : (0.98, 4.19)

approx 95% confidence interval for ψ : (0.98, 4.185)

$l_p(\hat{\psi}) - l_p(\psi)$

Evans et al. 1999

Logistic regression

The first ten out of 79 sets of observations on the physical characteristics of urine. Presence/absence of calcium oxalate crystals is indicated by 1/0. Two cases with missing values.

Case	Crystals	Specific gravity	pH	Osmolarity	Conductivity	Urea	Calcium
1	0	1.021	4.91	725	—	443	2.45
2	0	1.017	5.74	577	20.0	296	4.49
3	0	1.008	7.20	321	14.9	101	2.36
4	0	1.011	5.51	408	12.6	224	2.15
5	0	1.005	6.52	187	7.5	91	1.16
6	0	1.020	5.27	668	25.3	252	3.34
7	0	1.012	5.62	461	17.4	195	1.40
8	0	1.029	5.67	1107	35.9	550	8.48
9	0	1.015	5.41	543	21.9	170	1.16
10	0	1.021	6.13	779	25.7	382	2.21
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮

Andrews & Herzberg, 1985

Model: Independent binary responses Y_1, \dots, Y_n with

$$\Pr(Y_i = 1) = \frac{\exp(x_i^T \beta)}{1 + \exp(x_i^T \beta)}$$

Fitting generalized linear model in R:

```
data(urine)
fit <- glm(r~gravity+ph+osmo+cond+urea+calc,
family = binomial, data=urine)
summary(fit)
```

$$H_0: \beta_j = 0$$

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-355.33771	222.76696	-1.595	0.11069
gravity	355.94379	222.11004	1.603	0.10903
ph	-0.49570	0.56976	-0.870	0.38429
osmo	0.01681	0.01782	0.944	0.34536
conduct	-0.43282	0.25123	-1.723	0.08493 .
urea	-0.03201	0.01612	-1.986	0.04703 *
calc	0.78369	0.24216	3.236	0.00121 **

$$\hat{\beta}_j \approx \beta_j \quad s.e(\hat{\beta}_j)$$

$$\hat{\beta}_j / s.e(\hat{\beta}_j)$$

A closer look at coefficient of urea

method	lower bound	upper bound	p-value for 0
$\Phi(q)$	-0.063	-0.0006	0.047
$\Phi(r)$	-0.067	-0.0025	0.033
$\Phi(r^*)$	-0.058	0.0002	0.052

library(cond) # part of ~~package 'hoa'~~ on cran-r

glm's

```
urine.cond.urea <- cond.glm(urine.glm, offset=urea)
> summary(urine.cond.urea, test=0)
```

Brazzale

Test statistics


hypothesis : coef(urea) = 0

	statistic	tail prob.
Wald pivot	-1.986	0.02351
Wald pivot (cond. MLE)	-1.852	0.03202
Likelihood root	-2.133	0.01648
Modified likelihood root	-1.925	0.02713
Modified likelihood root (cont. corr.)	-1.917	0.02760

A closer look at coefficient of urea

method	lower bound	upper bound	p-value for 0
$\Phi(q)$	-0.063	-0.0006	0.047
$\Phi(r)$	-0.067	-0.0025	0.033
$\Phi(r^*)$	-0.058	0.0002	0.052

Profile and modified profile log likelihoods



Several 2×2 tables.

Institution	y_1	m_1	y_2	m_2	Institution	y_1	m_1	y_2	m_2
1	3	4	1	3	12	2	2	0	2
2	3	4	8	11	13	1	4	1	5
3	2	2	2	3	14	2	3	2	4
4	2	2	2	2	15	2	4	4	6
5	2	2	0	3	16	4	12	3	9
6	1	3	2	3	17	1	2	2	3
7	2	2	2	3	18	3	3	1	4
8	1	5	4	4	19	1	4	2	3
9	2	2	2	3	20	0	3	0	2
10	0	2	2	3	21	2	4	1	5
11	3	3	3	3					

Lipsitz et al. 1988: Biometrics

... matched pairs

Model: $Y_{1i} \sim \text{Binomial}(m_{1i}, p_{1i})$ $Y_{2i} \sim \text{Binomial}(m_{2i}, p_{2i})$

parameter of interest $\psi = p_{2i} - p_{1i}$

nuisance parameters $p_{1i}, i = 1, \dots, 21$

inference for ψ :

	lower	upper	point estimate	p -value for $\psi = 0$
$\Phi(r)$	-0.081	0.243	0.10	0.16
$\Phi(r^*)$	-0.137	0.126	0.01	0.45

... matched pairs

Model: $Y_{1i} \sim \text{Binomial}(m_{1i}, p_{1i})$ $Y_{2i} \sim \text{Binomial}(m_{2i}, p_{2i})$

parameter of interest $\psi = p_{2i} - p_{1i}$

nuisance parameters $p_{1i}, i = 1, \dots, 21$

inference for ψ :

	lower	upper	point estimate	p -value for $\psi = 0$
$\Phi(r)$	-0.081	0.243	0.10	0.16
$\Phi(r^*)$	-0.137	0.126	0.01	0.45

... matched pairs


Model: $Y_{1i} \sim \text{Binomial}(m_{1i}, p_{1i})$ $Y_{2i} \sim \text{Binomial}(m_{2i}, p_{2i})$

parameter of interest $\psi = p_{2i} - p_{1i}$


nuisance parameters $p_{1i}, i = 1, \dots, 21$

inference for ψ :

	lower	upper	point estimate	p -value for $\psi = 0$
$\Phi(r)$	-0.081	0.243	0.10	0.16
$\Phi(r^*)$	-0.137	0.126	0.01	0.45



likelihood root, r^* , q



profile log likelihood,
modified profile

Example G: Cox & Snell, 1980

	cost	date	T1	T2	cap	PR	NE	CT	BW	N	PT
1	460.05	68.58	14	46	687	0	1	0	0	14	0
2	452.99	67.33	10	73	1065	0	0	1	0	1	0
3	443.22	67.33	10	85	1065	1	0	1	0	1	0
4	652.32	68.00	11	67	1065	0	1	1	0	12	0
5	642.23	68.00	11	78	1065	1	1	1	0	12	0
6	345.39	67.92	13	51	514	0	1	1	0	3	0
7	272.37	68.17	12	50	822	0	0	0	0	5	0
8	317.21	68.42	14	59	457	0	0	0	0	1	0
9	457.12	68.42	15	55	822	1	0	0	0	5	0
10	690.19	68.33	12	71	792	0	1	1	1	2	0
11	350.63	68.58	12	64	560	0	0	0	0	3	0
12	402.59	68.75	13	47	790	0	1	0	0	6	0
13	412.18	68.42	15	62	530	0	0	1	0	2	0
14	495.58	68.92	17	52	1050	0	0	0	0	7	0
15	394.36	68.92	13	65	850	0	0	0	1	16	0

$$n = 32, d = 8$$

Linear regression, non-normal error

- Model $Y_i = \beta_0 + x_i^T \beta + \sigma \epsilon_i$
- $\epsilon \sim N(0, 1)$ or $\epsilon \sim t_\nu$

	Normal		t_4 , first order		t_4 , third order	
	Est (SE)	z	Est (SE)	z	Est (SE)	z
Constant	-13.26 (3.140)	-4.22	-11.30 (3.67)	-3.01	-11.86 (3.70)	-3.21
date	0.212 (0.043)	4.91	0.191 (0.048)	3.97	0.196 (0.049)	4.02
log(cap)	0.723 (0.119)	6.09	0.648 (0.113)	5.71	0.682 (0.129)	5.31
NE	0.249 (0.074)	3.36	0.242 (0.077)	3.12	0.239 (0.080)	2.97
CT	0.140 (0.060)	2.32	0.144 (0.054)	2.68	0.143 (0.063)	2.26
log(N)	-0.088 (0.042)	-2.11	-0.060 (0.043)	-1.40	-0.072 (0.048)	-1.51
PT	-0.226 (0.114)	-1.99	-0.282 (0.101)	-2.80	-0.265 (0.110)	-2.42

ν	log (N)		PT	
	First order	Third order	First order	Third order
4	0.162	0.151	0.005	0.024
6	0.110	0.116	0.007	0.032
8	0.081	0.098	0.009	0.036
10	0.064	0.086	0.011	0.038
20	0.036	0.064	0.016	0.045
40	0.025	0.053	0.029	0.050
100	0.020	0.047	0.022	0.053

Linear regression, non-normal error

- Model $Y_i = \beta_0 + x_i^T \beta + \sigma \epsilon_i$
- $\epsilon \sim N(0, 1)$ or $\epsilon \sim t_\nu$

	Normal		t_4 , first order		t_4 , third order	
	Est (SE)	z	Est (SE)	z	Est (SE)	z
Constant	-13.26 (3.140)	-4.22	-11.30 (3.67)	-3.01	-11.86 (3.70)	-3.21
date	0.212 (0.043)	4.91	0.191 (0.048)	3.97	0.196 (0.049)	4.02
log(cap)	0.723 (0.119)	6.09	0.648 (0.113)	5.71	0.682 (0.129)	5.31
NE	0.249 (0.074)	3.36	0.242 (0.077)	3.12	0.239 (0.080)	2.97
CT	0.140 (0.060)	2.32	0.144 (0.054)	2.68	0.143 (0.063)	2.26
log(N)	-0.088 (0.042)	-2.11	-0.060 (0.043)	-1.40	-0.072 (0.048)	-1.51
PT	-0.226 (0.114)	-1.99	-0.282 (0.101)	-2.80	-0.265 (0.110)	-2.42

ν	log (N)		PT	
	First order	Third order	First order	Third order
4	0.162	0.151	0.005	0.024
6	0.110	0.116	0.007	0.032
8	0.081	0.098	0.009	0.036
10	0.064	0.086	0.011	0.038
20	0.036	0.064	0.016	0.045
40	0.025	0.053	0.029	0.050
100	0.020	0.047	0.022	0.053
200	0.015	0.045	0.046	0.057

Linear regression, non-normal error

- Model $Y_i = \beta_0 + x_i^T \beta + \sigma \epsilon_i$
- $\epsilon \sim N(0, 1)$ or $\epsilon \sim t_\nu$

library (marg)

	Normal		t_4 , first order		t_4 , third order	
	Est (SE)	z	Est (SE)	z	Est (SE)	z
Constant	-13.26 (3.140)	-4.22	-11.30 (3.67)	-3.01	-11.86 (3.70)	-3.21
date	0.212 (0.043)	4.91	0.191 (0.048)	3.97	0.196 (0.049)	4.02
log(cap)	0.723 (0.119)	6.09	0.648 (0.113)	5.71	0.682 (0.129)	5.31
NE	0.249 (0.074)	3.36	0.242 (0.077)	3.12	0.239 (0.080)	2.97
CT	0.140 (0.060)	2.32	0.144 (0.054)	2.68	0.143 (0.063)	2.26
log(N)	-0.088 (0.042)	-2.11	-0.060 (0.043)	-1.40	-0.072 (0.048)	-1.51
PT	-0.226 (0.114)	-1.99	-0.282 (0.101)	-2.80	-0.265 (0.110)	-2.42

ν	log (N)		PT	
	First order	Third order	First order	Third order
4	0.162	0.151	0.005	0.024
6	0.110	0.116	0.007	0.032
8	0.081	0.098	0.009	0.036
10	0.064	0.086	0.011	0.038
20	0.036	0.064	0.016	0.045
40	0.025	0.053	0.029	0.050
100	0.020	0.047	0.022	0.053
2035	0.045	0.046	0.057	

Linear regression, non-normal error

- Model $Y_i = \beta_0 + x_i^T \beta + \sigma \epsilon_i$
- $\epsilon \sim N(0, 1)$ or $\epsilon \sim t_\nu$

	Normal		t_4 , first order		t_4 , third order	
	Est (SE)	z	Est (SE)	z	Est (SE)	z
Constant	-13.26 (3.140)	-4.22	-11.30 (3.67)	-3.01	-11.86 (3.70)	-3.21
date	0.212 (0.043)	4.91	0.191 (0.048)	3.97	0.196 (0.049)	4.02
log(cap)	0.723 (0.119)	6.09	0.648 (0.113)	5.71	0.682 (0.129)	5.31
NE	0.249 (0.074)	3.36	0.242 (0.077)	3.12	0.239 (0.080)	2.97
CT	0.140 (0.060)	2.32	0.144 (0.054)	2.68	0.143 (0.063)	2.26
log(N)	-0.088 (0.042)	-2.11	-0.060 (0.043)	-1.40	-0.072 (0.048)	-1.51
PT	-0.226 (0.114)	-1.99	-0.282 (0.101)	-2.80	-0.265 (0.110)	-2.42

ν	log (N)		PT	
	First order	Third order	First order	Third order
4	0.162	0.151	0.005	0.024
6	0.110	0.116	0.007	0.032
8	0.081	0.098	0.009	0.036
10	0.064	0.086	0.011	0.038
20	0.036	0.064	0.016	0.045
40	0.025	0.053	0.029	0.050
100	0.020	0.047	0.022	0.053
	0.035	0.045	0.046	0.057

```
library(marg)
# part of package 'hoa' on cran-r
data(nuclear)

# Fit normal-theory linear model and examine its contents:

nuc.norm <- lm( log(cost) ~ date + log(cap) + NE + CT + log(N) + PT,
+                 data = nuclear )
summary(nuc.norm)

# Fit linear model with t errors and 4 df and examine its contents:

nuc.t4 <- rsm( log(cost) ~ date + log(cap) + NE + CT + log(N) + PT,
+                 data = nuclear, family = student(4) )
summary(nuc.t4)
plot(nuc.t4)

# Conditional analysis for partial turnkey guarantee:

nuc.t4.pt <- cond( nuc.t4, offset = PT )
summary(nuc.t4.pt)
plot(nuc.t4.pt)

# For conditional analysis for other covariates, replace pt by
# log(N), ...
```

Type II censored data

40 units on test, 28 failures at (log) times

0.0507	0.0579	0.0784	0.0954	0.1376	0.2249	0.2362	0.2481
0.2501	0.2811	0.3027	0.3091	0.4296	0.5379	0.5621	0.5781
0.7811	0.8228	0.9455	0.9871	1.0060	1.0335	1.0377	1.0471
1.0876	1.2473	1.2776	1.3445				

Weibull model: $f(y; \mu, \sigma) = e^{(y-\mu)/\sigma} \exp\{-e^{(y-\mu)/\sigma}\}$

90% confidence intervals

	μ	σ
$\Phi(r)$	(-0.116, 0.476)	(0.700, 1.217)
$\Phi(r^*)$	(-0.107, 0.510)	(0.743, 1.320)
Exact (num. int.)	(-0.11, 0.51)	(0.724, 1.277)

Lawless 2003 Ch.5; Wong & Wu 2003

Type II censored data

40 units on test, 28 failures at (log) times

0.0507	0.0579	0.0784	0.0954	0.1376	0.2249	0.2362	0.2481
0.2501	0.2811	0.3027	0.3091	0.4296	0.5379	0.5621	0.5781
0.7811	0.8228	0.9455	0.9871	1.0060	1.0335	1.0377	1.0471
1.0876	1.2473	1.2776	1.3445				

Weibull model: $f(y; \mu, \sigma) = e^{(y-\mu)/\sigma} \exp\{-e^{(y-\mu)/\sigma}\}$

90% confidence intervals

	μ	σ
$\Phi(r)$	(-0.116, 0.476)	(0.700, 1.217)
$\Phi(r^*)$	(-0.107, 0.510)	(0.743, 1.320)
Exact (num. int.)	(-0.11, 0.51)	(0.724, 1.277)

Lawless 2003 Ch.5; Wong & Wu 2003

Type II censored data

40 units on test, 28 failures at (log) times

0.0507	0.0579	0.0784	0.0954	0.1376	0.2249	0.2362	0.2481
0.2501	0.2811	0.3027	0.3091	0.4296	0.5379	0.5621	0.5781
0.7811	0.8228	0.9455	0.9871	1.0060	1.0335	1.0377	1.0471
1.0876	1.2473	1.2776	1.3445				

Weibull model: $f(y; \mu, \sigma) = e^{(y-\mu)/\sigma} \exp\{-e^{(y-\mu)/\sigma}\}$

90% confidence intervals

	μ	σ
$\Phi(r)$	(-0.116, 0.476)	(0.700, 1.217)
$\Phi(r^*)$	(-0.107, 0.510)	(0.743, 1.320)
Exact (num. int.)	(-0.11, 0.51)	(0.724, 1.277)

Lawless 2003 Ch.5; Wong & Wu 2003

Type II censored data

40 units on test, 28 failures at (log) times

0.0507	0.0579	0.0784	0.0954	0.1376	0.2249	0.2362	0.2481
0.2501	0.2811	0.3027	0.3091	0.4296	0.5379	0.5621	0.5781
0.7811	0.8228	0.9455	0.9871	1.0060	1.0335	1.0377	1.0471
1.0876	1.2473	1.2776	1.3445				

Weibull model: $f(y; \mu, \sigma) = e^{(y-\mu)/\sigma} \exp\{-e^{(y-\mu)/\sigma}\}$

90% confidence intervals

	μ	σ
$\Phi(r)$	(-0.116, 0.476)	(0.700, 1.217)
$\Phi(r^*)$	(-0.107, 0.510)	(0.743, 1.320)
Exact (num. int.)	(-0.11, 0.51)	(0.724, 1.277)

Lawless 2003 Ch.5; Wong & Wu 2003

Vector parameter of interest

- ▶ use tangent exponential model (or usual exponential family model)
- ▶ construct a scalar parameter of interest representing direction in sample space
- ▶ apply higher order approximation
- ▶ Example $y \sim N(\mu, \Sigma)$; $H_0 : \Sigma^{-1}$ is tri-diagonal
- ▶ First-order Markov dependence in a graphical model

Nominal (%)	1.0	2.5	5.0	10.0	25.0	50.0	75.0	90.0	95.0	97.5
First order	5.5	10.5	17.0	27.0	48.7	73.0	89.5	96.7	98.5	99.4
Second order	1.1	2.6	5.0	10.1	24.8	49.8	74.9	89.9	94.9	97.4

